3625 Del Amo Boulevard, Suite 180 Torrance, California 90503-1643 (310) 370-8370 (310) 370-7026 FAX www.hygienetech.com

March 4, 2014

California State Board of Equalization 450 N Street Sacramento, California 94279

Document No. 21401001.4

Attention: David Gau

Regarding: Limited Fungal Growth Exposure Assessment Surveys

February 2014 Random Sampling

Dear Mr. Gau:

On February 7, 14, 20, and 24, 2014, industrial hygienists with Hygiene Technologies International, Inc. (HygieneTech) conducted limited fungal growth exposure assessment surveys involving 22 randomly selected areas located within the California State Board of Equalization (BOE) building. The findings of the surveys, along with the analytical data, conclusions, and recommendations when applicable, appear below.

On the survey dates, air samples were collected for total (viable and nonviable) fungi analyses using a Zefon brand Bio-Pump $Plus^{\mathsf{T}}$ equipped with Air-O-Cell cassettes. All such samples were subsequently analyzed for fungi (including yeasts, molds, rusts, smuts, and mushrooms) by trained and experienced microbiologists at a laboratory accredited by the American Industrial Hygiene Association (AIHA) and that successfully participates in the AIHA Environmental Microbiology Proficiency Analytical Testing (EMPAT) Program. The airborne fungi assessment analytical data with supporting and background information appear in the enclosed table.

As presented in Table 21402001-4, the airborne spore count data recorded showed fungal spore types outdoors such as *Alternaria*, ascospores, basidiospores, *Cladosporium*, colorless spores typical of *Penicillium/Aspergillus* species, *Curvularia*, *Epicoccum*, other brown, *Pithomyces*, rusts, smuts, *Stemphylium*, and/or *Torula*. In the indoor areas tested, the data showed that airborne fungal spores were either not detected at or above the laboratory detection limit indicated or were detected at low airborne concentrations. The fungal spore types found indoor included *Alternaria*, ascospores, basidiospores, *Cladosporium*, colorless spores typical of *Penicillium/Aspergillus* species, *Oidium*, other brown, rusts, and/or smuts. The distribution of fungal spore types detected in the surveyed areas was consistent with those found outdoors, and the overall data within the tested areas were well below the overall outdoor data recorded. These data are considered unremarkable and are not believed to pose a health risk beyond that posed by the outdoor environment where exposures to airborne fungi are expected.

Be advised that the data provided in this report only represent limited fungal growth and exposure potentials that existed at the time these surveys were performed and at the precise sample locations

Mr. David Gau March 4, 2014

Document No. 21402001.4 - February 2014 Random Sampling

Page 2



indicated. Note that fungal growth and exposure potentials may change due to changes in environmental conditions (such as those caused by water intrusion), use of mechanical systems, or other factors. Also be advised that additional fungal growth may exist at one or more locations in the structure that were not specifically assessed during the surveys.

If you have any comments or questions regarding the information contained in this correspondence, please feel free to contact our offices directly at (310) 370-8370.

Sincerely,

HYGIENE TECHNOLOGIES INTERNATIONAL, INC.

Kenny K. Hsi, CIH

Technical Director

APPENDIX A

CLIENT: California State Board of Equalization 450 N Street Sacramento, California 94279 TABLE 21402001-4
AIRBORNE TOTAL FUNGI RESULTS
450 N STREET
SACRAMENTO, CALIFORNIA
FEBRUARY 7, 14, 20, AND 24, 2014

Page 1

Results reported in spores per cubic meter of air (spores/M³)

Results reported in spores per cubic meter of air (spores/M ³)										
SAMPLE NUMBER	21402001-4 TM01OUT	21402001-4 TM02	21402001-4 TM03	21402001-4 TM04						
SAMPLING LOCATION/ACTIVITIES	Outdoors; southeast corner of building; approximately five feet above ground/Normal outdoor activities	20 th Floor; southern corridor; about two feet south of Mail Room 20B entry door; approximately five feet above floor/Normal office activities	18 th Floor; Break Room 1805; about center; approximately five feet above floor/Normal office activities	14 th Floor; Copy Room 1405; about center; approximately five feet above floor/Normal office activities						
DATE	02/07/14	02/07/14	02/07/14	02/07/14						
START/STOP	09:56:00/10:01:00	10:38:00/10:43:00	10:46:00/10:51:00	10:53:00/10:58:00						
SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes						
Alternaria	270									
Ascospores										
Basidiospores	1,000									
Bipolaris/Drechslera group										
Botrytis										
Chaetomium										
Cladosporium	3,000	110	53	160						
Curvularia										
Epicoccum										
Fusarium										
Nigrospora										
Oidium										
Other brown										
Other colorless										
Penicillium/Aspergillus types										
Pithomyces										
Rusts										
Smuts (Periconia, Myxomycetes)			13							
Stachybotrys										
Stemphylium										
Torula										
Ulocladium										
Hyphal fragments	<13	<13	13	<13						
Background debris*	1+	1+	2+	2+						
TOTAL**	4,300	110	67	160						

^{*}Background debris is an indication of the amount of non-biological particulate matter present on the slide and is graded (from least to greatest) as 1+ to 4+.

^{**}Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.



CLIENT: California State Board of Equalization 450 N Street Sacramento, California 94279

TABLE 21402001-4 **AIRBORNE TOTAL FUNGI RESULTS 450 N STREET** SACRAMENTO, CALIFORNIA FEBRUARY 7, 14, 20, AND 24, 2014

Page 2

Results reported in spores per cubic meter of air (spores/M³)										
SAMPLE NUMBER	21402001-4 TM05	21402001-4 TM06	21402001-4 TM07OUT	21402001-4 TM08						
SAMPLING LOCATION/ACTIVITIES	9 th Floor; Column N23 area; Cubicle 86 entry area; approximately five feet above floor/Normal office activities	4 th Floor; Column K17 area; Cubicle 76; immediately adjacent to Column K17; approximately five feet above floor/Normal office activities	Outdoors; about 15 feet east of building; approximately five feet above ground/Normal outdoor activities	1st Floor; Cafeteria; Café counter area; immediately east of counter; approximately five feet above floor/Normal office activities						
DATE	02/07/14	02/07/14	02/14/14	02/14/14						
START/STOP	11:02:00/11:07:00	11:11:00/11:16:00	15:25:00/15:30:00	15:33:00/15:38:00						
SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes						
Alternaria			27							
Ascospores			1,900							
Basidiospores	53		800							
Bipolaris/Drechslera group										
Botrytis										
Chaetomium										
Cladosporium	53		9,200							
Curvularia										
Epicoccum			40							
Fusarium										
Nigrospora										
Oidium										
Other brown				27						
Penicillium/Aspergillus types			480							
Pithomyces										
Rusts										
Smuts (Periconia, Myxomycetes)		13		13						
Stachybotrys										
Stemphylium			67							
Torula										
Trichocladium										
Ulocladium										
Zygomycetes										
Hyphal fragments	<13	<13	<13	<13						
Background debris*	2+	2+	1+	2+						
TOTAL**	110	13	13,000	40						

^{*}Background debris is an indication of the amount of non-biological particulate matter present on the slide and is graded (from least to greatest) as 1+ to 4+.

^{**}Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.



CLIENT: California State Board of Equalization 450 N Street Sacramento, California 94279

TABLE 21402001-4 AIRBORNE TOTAL FUNGI RESULTS **450 N STREET** SACRAMENTO, CALIFORNIA FEBRUARY 7, 14, 20, AND 24, 2014

Page 3

Results reported in spores per cubic meter of air (spores/M ³)										
SAMPLE NUMBER	21402001-4 TM09	21402001-4 TM10	21402001-4 TM11	21402001-4 TM12						
SAMPLING LOCATION/ACTIVITIES	3 rd Floor; Column O22 area; Cubicle at Column O22; approximately five feet above floor/Normal office activities	7 th Floor; Break Room 707; about center; approximately five feet above floor/Normal office activities	17 th Floor; Room 1708; about five feet southeast of entry door; approximately five feet above floor/Normal office activities	19 th Floor; Room 1908; about center; approximately five feet above floor/Normal office activities						
DATE	02/14/14	02/14/14	02/14/14	02/14/14						
START/STOP	15:42:00/15:47:00	15:51:00/15:56:00	15:59:00/16:04:00	16:06:00/16:11:00						
SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes						
Alternaria										
Arthrinium										
Ascospores				53						
Basidiospores				53						
Bipolaris/Drechslera group										
Botrytis										
Chaetomium										
Cladosporium	53	53		53						
Curvularia										
Epicoccum										
Fusarium										
Nigrospora										
Oidium										
Other brown		13								
Penicillium/Aspergillus types		53								
Pithomyces										
Rusts										
Smuts (Periconia, Myxomycetes)										
Stachybotrys										
Stemphylium										
Torula										
Ulocladium										
Hyphal fragments	<13	<13	<13	13						
Background debris*	2+	2+	1+	2+						
TOTAL**	53	120	<13	160						

^{*}Background debris is an indication of the amount of non-biological particulate matter present on the slide and is graded (from least to greatest) as 1+ to 4+.

^{**}Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.



CLIENT: California State Board of Equalization 450 N Street Sacramento, California 94279 TABLE 21402001-4
AIRBORNE TOTAL FUNGI RESULTS
450 N STREET
SACRAMENTO, CALIFORNIA
FEBRUARY 7, 14, 20, AND 24, 2014

Page 4

Results reported in spores per cubic meter of air (spores/M³)

Results reported in spores per cubic meter of air (spores/M ³)									
SAMPLE NUMBER	21402001-4 TM13	21402001-4 TM14OUT	21402001-4 TM15	21402001-4 TM16					
SAMPLING LOCATION/ACTIVITIES	22 nd Floor; Column K18 area; about 10 feet southwest of Column K18; approximately five feet above floor/Normal office activities	Outdoors; about 15 feet west of building; approximately five feet above ground/Normal outdoor activities	5 th Floor; Quiet Room 504; entry area; approximately five feet above floor/Normal office activities	8 th Floor; Room 809; about center; approximately five feet above floor/Normal office activities					
DATE	02/14/14	02/20/14	02/20/14	02/20/14					
START/STOP	16:15:00/16:20:00	14:49:00/14:54:00	14:58:00/15:03:00	15:05:00/15:10:00					
SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes					
Alternaria		67							
Ascospores		160							
Basidiospores		1,000	210	53					
Bipolaris/Drechslera group									
Botrytis									
Chaetomium									
Cladosporium		4,300	320	210					
Curvularia		13							
Epicoccum		13							
Myrothecium									
Nigrospora									
Oidium									
Other colorless									
Penicillium/Aspergillus types		320							
Pithomyces		13							
Rusts		13							
Smuts (Periconia, Myxomycetes)		40	13						
Stachybotrys									
Stemphylium									
Torula									
Ulocladium									
Zygomycetes									
Hyphal fragments	<13	93	40	<13					
Background debris*	1+	2+	2+	2+					
TOTAL**	<13	6,000	550	270					

^{*}Background debris is an indication of the amount of non-biological particulate matter present on the slide and is graded (from least to greatest) as 1+ to 4+.

^{**}Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.

APPENDIX A

CLIENT: California State Board of Equalization 450 N Street Sacramento, California 94279 TABLE 21402001-4
AIRBORNE TOTAL FUNGI RESULTS
450 N STREET
SACRAMENTO, CALIFORNIA
FEBRUARY 7, 14, 20, AND 24, 2014

Page 5

Results reported in spores per cubic meter of air (spores/M³)

Results reported in spores per cubic meter of air (spores/M ³)										
SAMPLE NUMBER	21402001-4 TM17	21402001-4 TM18	21402001-4 TM19	21402001-4 TM20OUT						
SAMPLING LOCATION/ACTIVITIES	15 th Floor; Break Room 1508; about center; approximately five feet above floor/Normal office	16 th Floor; Break Room 1603; about center; approximately five feet above floor/Normal office	24 th Floor; Room 2447; about center; approximately five feet above floor/Normal office activities	Outdoors; about 10 feet south of building; approximately five feet above ground/Normal						
	activities	activities	Office activities	outdoor activities						
DATE	02/20/14	02/20/14	02/20/14	02/24/14						
START/STOP	15:13:00/15:18:00	15:20:00/15:25:00	15:28:00/15:33:00	14:49:00/14:54:00						
SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes						
Alternaria		13		40						
Ascospores				590						
Basidiospores	53	210	53	1,300						
Bipolaris/Drechslera group										
Botrytis										
Chaetomium										
Cladosporium	53	370	53	4,800						
Curvularia										
Epicoccum										
Fusarium										
Myrothecium										
Nigrospora										
Oidium										
Other brown				13						
Penicillium/Aspergillus types			110	110						
Pithomyces										
Rusts				53						
Smuts (Periconia, Myxomycetes)				80						
Stemphylium				13						
Stachybotrys										
Torula				80						
Ulocladium										
Zygomycetes										
Hyphal fragments	13	27	<13	53						
Background debris*	2+	2+	2+	2+						
TOTAL**	110	600	210	7,100						

^{*}Background debris is an indication of the amount of non-biological particulate matter present on the slide and is graded (from least to greatest) as 1+ to 4+.

^{**}Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.



CLIENT: California State Board of Equalization 450 N Street Sacramento, California 94279 TABLE 21402001-4
AIRBORNE TOTAL FUNGI RESULTS
450 N STREET
SACRAMENTO, CALIFORNIA
FEBRUARY 7, 14, 20, AND 24, 2014

Page 6

Results reported in spores per cubic meter of air (spores/M³)

Results reported in spores per cubic meter of air (spores/M³)										
SAMPLE NUMBER	21402001-4 TM21	21402001-4 TM22	21402001-4 TM23	21402001-4 TM24						
SAMPLING LOCATION/ACTIVITIES	2 nd Floor; Column M18 area; about two feet north of Column M18; approximately five feet above floor/Normal office activities	6 th Floor; Column N18 area; about two feet southeast of Column N18; approximately five feet above floor/Normal office activities	10 th Floor; southern and western corridor junction; approximately feet above floor/Normal office activities	11 th Floor; Column J21 area; Cubicle 11 entry area; approximately five feet above floor/Normal office activities						
DATE	02/24/14	02/24/14	02/24/14	02/24/14						
START/STOP	14:59:00/15:04:00	15:10:00/15:15:00	15:17:00/15:22:00	15:25:00/15:30:00						
SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes						
Alternaria										
Ascospores										
Basidiospores			210							
Bipolaris/Drechslera group										
Botrytis										
Chaetomium										
Cladosporium			110	53						
Curvularia										
Epicoccum										
Fusarium										
Myrothecium										
Nigrospora										
Oidium				13						
Other brown			13	13						
Penicillium/Aspergillus types										
Pithomyces										
Rusts				13						
Smuts (Periconia, Myxomycetes)										
Stachybotrys										
Stemphylium										
Torula										
Ulocladium										
Hyphal fragments	<13	<13	<13	<13						
Background debris*	2+	2+	2+	2+						
TOTAL**	<13	<13	330	93						

^{*}Background debris is an indication of the amount of non-biological particulate matter present on the slide and is graded (from least to greatest) as 1+ to 4+.

^{**}Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.

APPENDIX A

CLIENT: California State Board of Equalization 450 N Street Sacramento, California 94279 TABLE 21402001-4
AIRBORNE TOTAL FUNGI RESULTS
450 N STREET
SACRAMENTO, CALIFORNIA
FEBRUARY 7, 14, 20, AND 24, 2014

				Page 7
SAMPLE NUMBER	21402001-4 TM25	21402001-4 TM26		
SAMPLING LOCATION/ACTIVITIES	21 st Floor; Mail Room 21B; about six feet north of entry door; approximately five feet	23 rd Floor; Room 2311; Reception area; about center; approximately five feet above	This column intentionally left blank	This column intentionally left blank
	above floor/Normal office activities	floor/Normal office activities		
DATE	02/24/14	02/24/14		
START/STOP	15:36:00/15:41:00	15:45:00/15:50:00		
SAMPLE TIME	5 minutes	5 minutes		
Alternaria				
Ascospores				
Basidiospores	53			
Bipolaris/Drechslera group				
Botrytis				
Chaetomium				
Cladosporium	110	53		
Curvularia				
Epicoccum				
Fusarium				
Myrothecium				
Nigrospora				
Oidium				
Other brown				
Penicillium/Aspergillus types		53		
Pithomyces				
Rusts	13			
Smuts (Periconia, Myxomycetes)				
Stachybotrys				
Stemphylium				
Torula				
Ulocladium				
Hyphal fragments	<13	<13		
Background debris*	2+	2+		
TOTAL**	170	110		

^{*}Background debris is an indication of the amount of non-biological particulate matter present on the slide and is graded (from least to greatest) as 1+ to 4+.

^{**}Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.



Report for:

Mr. Lakhpreet Sandhu Hygiene Technologies International, Inc. 3625 Del Amo Boulevard, Suite 180 Torrance, CA 90503-8370

Regarding: Project: 21402001-4

EMĹ ID: 1168914

Approved by:

Dates of Analysis: Spore trap analysis: 02-10-2014

Technical Manager Melissa Tracey

Service SOPs: Spore trap analysis (1038) AIHA-LAP, LLC accredited service, Lab ID #179768

All samples were received in acceptable condition unless noted in the Report Comments portion in the body of the report. Due to the nature of the analyses performed, field blank correction of results is not applied. The results relate only to the items tested.

EMLab P&K ("the Company") shall have no liability to the client or the client's customer with respect to decisions or recommendations made, actions taken or courses of conduct implemented by either the client or the client's customer as a result of or based upon the Test Results. In no event shall the Company be liable to the client with respect to the Test Results except for the Company's own willful misconduct or gross negligence nor shall the Company be liable for incidental or consequential damages or lost profits or revenues to the fullest extent such liability may be disclaimed by law, even if the Company has been advised of the possibility of such damages, lost profits or lost revenues. In no event shall the Company's liability with respect to the Test Results exceed the amount paid to the Company by the client therefor.

Date of Sampling: 02-07-2014

Client: Hygiene Technologies International, Inc.

Date of Receipt: 02-07-2014 C/O: Mr. Lakhpreet Sandhu Re: 21402001-4 Date of Report: 02-10-2014

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:	21402001-	-4 TM01OUT	2140200	01-4 TM02	21402001-4 TM03	
Comments (see below)	None		None		1	None
Lab ID-Version‡:	528	9805-1	5289806-1		5289807-1	
Analysis Date:	02/1	10/2014	02/1	0/2014	02/1	10/2014
	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3
Ascospores	5	270		_		_
Basidiospores	19	1,000				
Bipolaris/Drechslera group		,				
Botrytis						
Chaetomium						
Cladosporium	57	3,000	2	110	1	53
Curvularia		,				
Epicoccum						
Fusarium						
Myrothecium						
Nigrospora						
Other colorless						
Penicillium/Aspergillus types†						
Pithomyces						
Rusts						
Smuts, Periconia, Myxomycetes					1	13
Stachybotrys						
Stemphylium						
Torula						
Ulocladium						
Zygomycetes						
Background debris (1-4+)††	1+		1+		2+	
Hyphal fragments/m3	< 13		< 13		13	
Pollen/m3	67		13		< 13	
Skin cells (1-4+)	< 1+		< 1+		1+	
Sample volume (liters)	75		75		75	
§ TOTAL SPORES/m3		4,300		110		67

Comments:

Spore types listed without a count or data entry were not detected during the course of the analysis for the respective sample.

The analytical sensitivity is the spores/m3 divided by the raw count. The limit of detection is the analytical sensitivity multiplied by the sample volume divided by 1000.

EMLab P&K, LLC

EMLab ID: 1168914, Page 2 of 3

[†] The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††}Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be higher then reported. It is important to account for samples volumes when evaluating dust levels.

For more information regarding analytical sensitivity, please contact QA by calling the laboratory. ‡ A "Version" indicated by -"x" after the Lab ID# with a value greater than 1 indicates a sample with amended data. The revision number is reflected by the value of "x".

[§] Total Spores/m3 has been rounded to two significant figures to reflect analytical precision.

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu Re: 21402001-4

Date of Sampling: 02-07-2014 Date of Receipt: 02-07-2014 Date of Report: 02-10-2014

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:	214020	01-4 TM04	2140200	01-4 TM05	21402001-4 TM06	
Comments (see below)	- N	None	None		1	None
Lab ID-Version‡:	528	9808-1	5289809-1		5289810-1	
Analysis Date:	02/1	0/2014	02/1	0/2014	02/10/2014	
	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3
Ascospores		_		-		
Basidiospores			1	53		
Bipolaris/Drechslera group						
Botrytis						
Chaetomium						
Cladosporium	3	160	1	53		
Curvularia						
Epicoccum						
Fusarium						
Myrothecium						
Nigrospora						
Other colorless						
Penicillium/Aspergillus types†						
Pithomyces						
Rusts						
Smuts, Periconia, Myxomycetes					1	13
Stachybotrys						
Stemphylium						
Torula						
Ulocladium						
Zygomycetes						
Background debris (1-4+)††	2+		2+		2+	
Hyphal fragments/m3	< 13		< 13		< 13	
Pollen/m3	< 13		< 13		< 13	
Skin cells (1-4+)	1+		1+		1+	
Sample volume (liters)	75		75		75	
§ TOTAL SPORES/m3		160		110		13

Comments:

Spore types listed without a count or data entry were not detected during the course of the analysis for the respective sample.

The analytical sensitivity is the spores/m3 divided by the raw count. The limit of detection is the analytical sensitivity multiplied by the sample volume divided by 1000.

EMLab P&K, LLC

EMLab ID: 1168914, Page 3 of 3

[†] The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††}Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be higher then reported. It is important to account for samples volumes when evaluating dust levels.

For more information regarding analytical sensitivity, please contact QA by calling the laboratory. ‡ A "Version" indicated by -"x" after the Lab ID# with a value greater than 1 indicates a sample with amended data. The revision number is reflected by the value of "x".

[§] Total Spores/m3 has been rounded to two significant figures to reflect analytical precision.

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-07-2014 Date of Receipt: 02-07-2014 Date of Report: 02-10-2014

MoldRANGETM: Extended Outdoor Comparison

Outdoor Location: 21402001-4 TM01OUT

Fungi Identified	Outdoor		Typica	l Outo	loor Da	ta for	:	,	Typica	l Outo	loor Da	ata for	:
	data	Feb	ruary ir	n Calif	ornia† (n‡=16	302)	The entire year in California† (n‡=199769)					
	spores/m3	very low	low	med	high	very high	freq %	very low	low	med	high	very high	freq %
Generally able to grow indoors*													
Alternaria	-	13	13	20	44	67	38	13	13	27	67	110	54
Bipolaris/Drechslera group	-	7	13	13	27	40	7	7	13	13	27	40	12
Chaetomium	-	7	13	13	27	40	9	8	13	13	27	47	19
Cladosporium	3,000	80	150	400	1,000	1,700	95	110	210	610	1,600	2,800	97
Curvularia	-	7	10	13	13	27	2	7	13	13	27	53	6
Nigrospora	-	7	13	13	13	27	4	7	13	13	27	53	8
Penicillium/Aspergillus types	-	53	66	200	490	820	82	53	100	210	590	1,000	84
Stachybotrys	-	13	13	13	40	80	3	7	13	13	33	67	4
Torula	-	7	13	13	40	53	5	8	13	13	40	67	12
Seldom found growing indoors**													
Ascospores	270	27	53	160	510	960	72	25	53	110	360	690	71
Basidiospores	1,000	53	110	430	1,800	3,900	95	53	80	270	990	2,300	93
Rusts	-	8	13	13	40	73	14	13	13	13	53	80	26
Smuts, Periconia, Myxomycetes	-	13	13	27	67	110	54	13	13	40	110	210	68
§ TOTAL SPORES/m3	4,300												

[†]The "Typical Outdoor Data' represents the typical outdoor spore levels for the location and time frame indicated. The last column represents the frequency of occurrence. The very low, low, med, high, and very high values represent the 10, 20, 50, 80, and 90 percentile values of the spore type when it is detected. For example, if the frequency of occurrence is 63% and the low value is 53, it would mean that the given spore type is detected 63% of the time and, when detected, 20% of the time it is present in levels above the detection limit and below 53 spores/m3. These values are updated periodically, and if enough data is not available to make a statistically meaningful assessment, it is indicated with a dash.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor data" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. In addition, EMLab P&K may not have received and tested a representative number of samples for every region or time period. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the use or interpretation of the data contained in, or any actions taken or omitted in reliance upon, this report.

EMLab P&K, LLC EMLab ID: 1168914, Page 1 of 1

[§] Total Spores/m3 has been rounded to two significant figures to reflect analytical precision.

^{*} The spores in this category are generally capable of growing on wet building materials in addition to growing outdoors. Building related growth is dependent upon the fungal type, moisture level, type of material, and other factors. *Cladosporium* is one of the predominant spore types worldwide and is frequently present in high numbers. *Penicillium/Aspergillus* species colonize both outdoor and indoor wet surfaces rapidly and are very easily dispersed. Other genera are usually present in lesser numbers.

^{**} These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens. However, in each group there are notable exceptions. For example, agents of wood decay are members of the basidiomycetes and high counts of a single morphological type of basidiospore on an inside sample should be considered significant.

 $[\]ddagger$ n = number of samples used to calculate data.

Date of Sampling: 02-07-2014

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Date of Receipt: 02-07-2014 Re: 21402001-4 Date of Report: 02-10-2014

MoldSTATTM: Supplementary Statistical Spore Trap Report

Outdoor Summary: 21402001-4 TM01OUT:

Species detected		Outdoor	r sample sj	ores/m3	Typical outdoor ranges	Freq.
	<100	1K	10K	>100K	(North America)	%
Ascospores				270	13 - 210 - 5,700	76
Basidiospores				1,000] 15 - 450 - 24,000	92
Cladosporium				3,000] 27 - 480 - 10,000	90
Penicillium/Aspergillus types				< 13] 13 - 170 - 2,700	68
Smuts, Periconia, Myxomycetes				< 13] 7 - 53 - 930	64
Total				4,300]	

The "Typical outdoor ranges" and "Freq. %" columns show the typical low, medium, and high spore counts per cubic meter and the frequency of occurrence for the given spore type. The low, medium, and high values represent the 2.5, 50, and 97.5 percentile values when the spore type is detected. For example, if the low value is 53 and the frequency of occurrence is 63%, it would mean that we typically detect the given spore type on 63 percent of all outdoor samples and, when detected, 2.5% of the time it is present in levels below 53 spores/m3.

Indoor Samples

Location: 21402001-4 TM02

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: 2%	dF: 4 Result: 0.2667 Critical value: 9.4877 Inside Similar: Yes	Resul	t: 0.5000	dF: 3 Result: 0.8750 Critical value: N/A Outside Similar: N/A	Score: 102 Result: Low
Species 1	Detected			Spores/m3	
		<100	1K	10K	>100K
	Cladosporium				110
	Total				110

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)			
Result: 1%	dF: 4 Result: 0.2667 Critical value: 9.4877 Inside Similar: Yes	Result: 0.4000	dF: 4 Result: 0.3500 Critical value: N/A Outside Similar: N/A	Score: 103 Result: Low			
Species	Detected		Spores/m3				
		<100 1K	10K	>100K			
	Cladosporium			53			
Smuts, Periconia, Myxomycetes				13			
	Total			67			

Date of Sampling: 02-07-2014

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Date of Receipt: 02-07-2014 Re: 21402001-4 Date of Report: 02-10-2014

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 21402001-4 TM04

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)				
Result: 3%	dF: 4 Result: 0.2667 Critical value: 9.4877 Inside Similar: Yes	Result: 0.5000	dF: 3 Result: 0.8750 Critical value: N/A Outside Similar: N/A	Score: 103 Result: Low				
Species 1	Detected	Spores/m3						
		<100 1K	10K	>100K				
	Cladosporium			160				
	Total			160				

Location: 21402001-4 TM05

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)			
Result: 2%	dF: 4 Result: 0.2667 Critical value: 9.4877 Inside Similar: Yes	Result: 0.8000	dF: 3 Result: 0.8750 Critical value: N/A Outside Similar: N/A	Score: 103 Result: Low			
Species 1	Detected		Spores/m3				
		<100 1K	10K	>100K			
	Basidiospores			53			
	Cladosporium			53			
	Total			110			

Location: 21402001-4 TM06

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)				
Result: < 1%	dF: 4 Result: 0.2667 Critical value: 9.4877 Inside Similar: Yes	Result: 0.0000	dF: 4 Result: -0.4000 Critical value: N/A Outside Similar: N/A	Score: 103 Result: Low				
Species 1	Detected	Spores/m3						
		<100 1K	10K	>100K				
Smuts, P	Periconia, Myxomycetes			13				
	Total			13				

^{*} The Friedman chi-square statistic is a non-parametric test that examines variation in a set of data (in this case, all indoor spore counts). The null hypothesis (H0) being tested is that there is no meaningful difference in the data for all indoor locations. The alternative hypothesis (used if the test disproves the null hypothesis) is that there is a difference between the indoor locations. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

EMLab P&K, LLC EMLab ID: 1168914, Page 2 of 3

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-07-2014 Date of Receipt: 02-07-2014 Date of Report: 02-10-2014

MoldSTATTM: Supplementary Statistical Spore Trap Report

** An agreement ratio is a simple method for assessing the similarity of two samples (in this case the indoor sample and the outdoor summary) based on the spore types present. A score of one indicates that the types detected in one location are the same as that in the other. A score of zero indicates that none of the types detected indoors are present outdoors. Typically, an agreement of 0.8 or higher is considered high.

*** The Spearman rank correlation is a non-parametric test that examines correlation between two sets of data (in this case the indoor location and the outdoor summary). The null hypothesis (H0) being tested is that the indoor and outdoor samples are unrelated. The alternative hypothesis (used if the test disproves the null hypothesis) is that the samples are similar. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

**** MoldSCORETM is a specialized method for examining air sampling data. It is a score between 100 and 300, with 100 indicating a greater likelihood that the airborne indoor spores originated from the outside, and 300 indicating a greater likelihood that they originated from an inside source. The Result displayed is based on the numeric score given and will be either Low, Medium, or High, indicating a low, medium, or high likelihood that the spores detected originated from an indoor source. EMLab P&Kreserves the right to, and may at anytime, modify or change the MoldScore algorithm without notice.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor ranges" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. With the statistical analysis provided, as with all statistical comparisons and analyses, false-positive and false-negative results can and do occur. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the data contained in, or any actions taken or omitted in reliance upon, this report.

EMLab P&K, LLC EMLab ID: 1168914, Page 3 of 3

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-07-2014 Date of Receipt: 02-07-2014 Date of Report: 02-10-2014

MoldSCORETM: **Spore Trap Report Outdoor Sample:** 21402001-4 TM01OUT

Fungi Identified	Oı	ıtdo	or	san	np	le	spoi	res	/m3	Raw	Spores/
_	<10	0		1K			10K	>	>100	count	m3
Generally able to grow indoors*											
Alternaria										ND	< 13
Bipolaris/Drechslera group										ND	< 13
Chaetomium										ND	< 13
Cladosporium										57	3,000
Curvularia										ND	< 13
Nigrospora										ND	< 13
Penicillium/Aspergillus types†										ND	< 13
Stachybotrys										ND	< 13
Torula										ND	< 13
Seldom found growing indoors**											
Ascospores										5	270
Basidiospores										19	1,000
Rusts										ND	< 13
Smuts, Periconia, Myxomycetes										ND	< 13
Total											4,320

Fungi Identified	Ind	oor	sam	ple	spor	es/n	13	Raw	Spores/
	<100		1 K		10K	>	100K	count	m3
Generally able to grow indoors*									
Alternaria								ND	< 13
Bipolaris/Drechslera group								ND	< 13
Chaetomium								ND	< 13
Cladosporium								2	110
Curvularia								ND	< 13
Nigrospora								ND	< 13
Penicillium/Aspergillus types†								ND	< 13
Stachybotrys								ND	< 13
Torula								ND	< 13
Seldom found growing indoors**									
Ascospores								ND	< 13
Basidiospores								ND	< 13
Rusts								ND	< 13
Smuts, Periconia, Myxomycetes								ND	< 13
Total									107

100	MoldSCORE; 100 200 300 Score								
			100						
			100						
			100						
			102						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
Final	Final MoldSCORE								

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-07-2014 Date of Receipt: 02-07-2014 Date of Report: 02-10-2014

MoldSCORETM: Spore Trap Report

Location: 21402001-4 TM03

Fungi Identified	Indo	or	samj	ole s	spore	s/m	13	Raw	Spores/
	<100		1K		10K	>1	00K	count	m3
Generally able to grow indoors*									
Alternaria								ND	< 13
Bipolaris/Drechslera group								ND	< 13
Chaetomium								ND	< 13
Cladosporium								1	53
Curvularia								ND	< 13
Nigrospora								ND	< 13
Penicillium/Aspergillus types†								ND	< 13
Stachybotrys								ND	< 13
Torula								ND	< 13
Seldom found growing indoors**									
Ascospores								ND	< 13
Basidiospores								ND	< 13
Rusts								ND	< 13
Smuts, Periconia, Myxomycetes								1	13
Total			•						67

	MoldSCORE:								
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			103						
Final Mo	ldSCO	RE	103						

Fungi Identified	Indoor sample spores/n								/m3 Raw			Spores/
	<100			K			10K		>10	0K	count	m3
Generally able to grow indoors*												
Alternaria											ND	< 13
Bipolaris/Drechslera group					Ш						ND	< 13
Chaetomium											ND	< 13
Cladosporium											3	160
Curvularia											ND	< 13
Nigrospora											ND	< 13
Penicillium/Aspergillus types†											ND	< 13
Stachybotrys											ND	< 13
Torula											ND	< 13
Seldom found growing indoors**												
Ascospores											ND	< 13
Basidiospores											ND	< 13
Rusts											ND	< 13
Smuts, Periconia, Myxomycetes					П						ND	< 13
Total												160

100 I	MoldSCORE; 200 300 Score								
			100						
			100						
			100						
			103						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
Final	Final MoldSCORE								

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-07-2014 Date of Receipt: 02-07-2014 Date of Report: 02-10-2014

MoldSCORETM: Spore Trap Report

Location: 21402001-4 TM05

Fungi Identified	Indo	or	sam	ple :	spore	es/n	13	Raw	Spores/
	<100		1K		10K	>1	100K	count	m3
Generally able to grow indoors*									
Alternaria								ND	< 13
Bipolaris/Drechslera group								ND	< 13
Chaetomium								ND	< 13
Cladosporium								1	53
Curvularia								ND	< 13
Nigrospora								ND	< 13
Penicillium/Aspergillus types†								ND	< 13
Stachybotrys								ND	< 13
Torula								ND	< 13
Seldom found growing indoors**									
Ascospores								ND	< 13
Basidiospores								1	53
Rusts								ND	< 13
Smuts, Periconia, Myxomycetes								ND	< 13
Total									107

1							
	MoldSC						
100	100 200 300						
			100				
			100				
			100				
			100				
			100				
			100				
			100				
			100				
			100				
			100				
			103				
			100				
			100				
Fina	Final MoldSCORE						

Fungi Identified	Inc	Indoor sample spores/m3							3	Raw	Spores/	
	<100			K			10K		>10	0K	count	m3
Generally able to grow indoors*												
Alternaria											ND	< 13
Bipolaris/Drechslera group											ND	< 13
Chaetomium											ND	< 13
Cladosporium											ND	< 13
Curvularia											ND	< 13
Nigrospora											ND	< 13
Penicillium/Aspergillus types†											ND	< 13
Stachybotrys											ND	< 13
Torula											ND	< 13
Seldom found growing indoors**												
Ascospores											ND	< 13
Basidiospores											ND	< 13
Rusts											ND	< 13
Smuts, Periconia, Myxomycetes											1	13
Total												13

100 N	MoldSCORE‡ 200 300 Score									
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			103							
Final 1	Final MoldSCORE									

Client: Hygiene Technologies International, Inc. C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-07-2014 Date of Receipt: 02-07-2014 Date of Report: 02-10-2014

MoldSCORETM: Spore Trap Report

* The spores in this category are generally capable of growing on wet building materials in addition to growing outdoors. Building related growth is dependent upon the fungal type, moisture level, type of material, and other factors. *Cladosporium* is one of the predominant spore types worldwide and is frequently present in high numbers. *Penicillium/Aspergillus* species colonize both outdoor and indoor wet surfaces rapidly and are very easily dispersed. Other genera are usually present in lesser numbers.

** These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens.

** These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens. However, in each group there are notable exceptions. For example, agents of wood decay are members of the basidiomycetes and high counts of

a single morphological type of basidiospore on an inside sample should be considered significant.

†The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods.

‡Rated on a scale from 100 to 300. A rating less than 150 is low and indicates a low probability of spores originating inside. A rating greater than 250 is high and indicates a high probability that the spores originated from inside, presumably from indoor mold growth. A rating between 150 and 250 indicates a moderate likelihood of indoor fungal growth. MoldSCORE is NOT intended for wall cavity samples. It is intended for ambient air samples in residences. Using the analysis on other samples (like wall cavity samples) will lead to misleading results.

EMLab P&K, LLC EMLab ID: 1168914, Page 4 of 4



Report for:

Mr. Lakhpreet Sandhu Hygiene Technologies International, Inc. 3625 Del Amo Boulevard, Suite 180 Torrance, CA 90503-8370

Regarding: Project: 21402001-4

EMĹ ID: 1172421

Approved by:

Dates of Analysis:

Spore trap analysis: 02-19-2014

Technical Manager Melissa Tracey

Service SOPs: Spore trap analysis (1038) AIHA-LAP, LLC accredited service, Lab ID #179768

All samples were received in acceptable condition unless noted in the Report Comments portion in the body of the report. Due to the nature of the analyses performed, field blank correction of results is not applied. The results relate only to the items tested.

EMLab P&K ("the Company") shall have no liability to the client or the client's customer with respect to decisions or recommendations made, actions taken or courses of conduct implemented by either the client or the client's customer as a result of or based upon the Test Results. In no event shall the Company be liable to the client with respect to the Test Results except for the Company's own willful misconduct or gross negligence nor shall the Company be liable for incidental or consequential damages or lost profits or revenues to the fullest extent such liability may be disclaimed by law, even if the Company has been advised of the possibility of such damages, lost profits or lost revenues. In no event shall the Company's liability with respect to the Test Results exceed the amount paid to the Company by the client therefor.

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-14-2014 Date of Receipt: 02-18-2014 Date of Report: 02-19-2014

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:		02001- 07OUT	214020	01-4TM08	214020	01-4TM09	214020	01-4TM10
Comments (see below)	N	lone	N	lone	N	Vone	N	Vone
Lab ID-Version‡:	530	7040-1	530	7041-1	530	7042-1	530	7043-1
Analysis Date:	02/1	9/2014	02/1	9/2014	02/1	9/2014	02/19/2014	
·	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3
Alternaria	2	27				-		-
Ascospores	36	1,900						
Basidiospores	15	800						
Chaetomium								
Cladosporium	173	9,200			1	53	1	53
Epicoccum	3	40						
Fusarium								
Myrothecium								
Nigrospora								
Oidium	3	40						
Other brown			2	27			1	13
Other colorless								
Penicillium/Aspergillus types†	9	480					1	53
Pithomyces								
Rusts								
Smuts, Periconia, Myxomycetes			1	13				
Stachybotrys								
Stemphylium	5	67						
Torula								
Ulocladium								
Zygomycetes								
Background debris (1-4+)††	1+		2+		2+		2+	
Hyphal fragments/m3	< 13		< 13		< 13		< 13	
Pollen/m3	170		13		< 13		< 13	
Skin cells (1-4+)	< 1+		1+		1+		1+	
Sample volume (liters)	75		75		75		75	
§ TOTAL SPORES/m3		13,000		40		53		120

Comments:

Spore types listed without a count or data entry were not detected during the course of the analysis for the respective sample.

The analytical sensitivity is the spores/m3 divided by the raw count. The limit of detection is the analytical sensitivity multiplied by the sample volume divided by 1000.

EMLab P&K, LLC EMLab ID: 1172421, Page 2 of 3

[†] The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††}Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be higher then reported. It is important to account for samples volumes when evaluating dust levels.

For more information regarding analytical sensitivity, please contact QA by calling the laboratory. ‡ A "Version" indicated by -"x" after the Lab ID# with a value greater than 1 indicates a sample with amended data. The revision number is reflected by the value of "x".

[§] Total Spores/m3 has been rounded to two significant figures to reflect analytical precision.

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu Re: 21402001-4

Date of Sampling: 02-14-2014 Date of Receipt: 02-18-2014 Date of Report: 02-19-2014

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:	214020	01-4TM11	214020	01-4TM12	214020	01-4TM13
Comments (see below)	1	None	N	Vone	1	None
Lab ID-Version‡:	530	7044-1	530	7045-1	530	7046-1
Analysis Date:	02/1	19/2014	02/1	9/2014	02/1	19/2014
	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3
Alternaria		_		_		_
Ascospores			1	53		
Basidiospores			1	53		
Chaetomium						
Cladosporium			1	53		
Curvularia						
Epicoccum						
Fusarium						
Myrothecium						
Nigrospora						
Oidium						
Other brown						
Other colorless						
Penicillium/Aspergillus types†						
Pithomyces						
Rusts						
Smuts, Periconia, Myxomycetes						
Stachybotrys						
Stemphylium						
Torula						
Ulocladium						
Zygomycetes						
Background debris (1-4+)††	1+		2+		1+	
Hyphal fragments/m3	< 13		13		< 13	
Pollen/m3	< 13		< 13		< 13	
Skin cells (1-4+)	< 1+		1+		< 1+	
Sample volume (liters)	75		75		75	
§ TOTAL SPORES/m3		< 13		160		< 13

Comments:

Spore types listed without a count or data entry were not detected during the course of the analysis for the respective sample.

The analytical sensitivity is the spores/m3 divided by the raw count. The limit of detection is the analytical sensitivity multiplied by the sample volume divided by 1000.

EMLab P&K, LLC EMLab ID: 1172421, Page 3 of 3

[†] The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††}Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be higher then reported. It is important to account for samples volumes when evaluating dust levels.

For more information regarding analytical sensitivity, please contact QA by calling the laboratory. ‡ A "Version" indicated by -"x" after the Lab ID# with a value greater than 1 indicates a sample with amended data. The revision number is reflected by the value of "x".

[§] Total Spores/m3 has been rounded to two significant figures to reflect analytical precision.

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-14-2014 Date of Receipt: 02-18-2014 Date of Report: 02-19-2014

MoldRANGETM: Extended Outdoor Comparison

Outdoor Location: 21402001-4TM07OUT

Fungi Identified	Outdoor		Typica	l Outo	loor Da	ıta for	:	1	Typica	l Outo	loor Da	ata for	:
	data	Feb	ruary ir	n Calif	ornia† (n‡=16	302)	The er	ntire yea	ar in Ca	lifornia	(n‡=2	00710)
	spores/m3	very low	low	med	high	very high	freq %	very low	low	med	high	very high	freq %
Generally able to grow indoors*													
Alternaria	27	13	13	20	44	67	38	13	13	27	67	110	53
Bipolaris/Drechslera group	-	7	13	13	27	40	7	7	13	13	27	40	12
Chaetomium	-	7	13	13	27	40	9	8	13	13	27	47	19
Cladosporium	9,200	80	150	400	1,000	1,700	95	110	210	610	1,600	2,800	97
Curvularia	-	7	10	13	13	27	2	7	13	13	27	53	6
Epicoccum	40	7	13	13	27	53	13	8	13	13	33	53	19
Nigrospora	-	7	13	13	13	27	4	7	13	13	27	53	8
Other brown	-	11	13	13	27	47	29	13	13	13	40	53	34
Penicillium/Aspergillus types	480	53	66	200	490	820	82	53	100	210	590	1,000	84
Stachybotrys	-	13	13	13	40	80	3	7	13	13	33	67	4
Stemphylium	67	7	13	13	27	40	5	7	13	13	27	40	9
Torula	-	7	13	13	40	53	5	8	13	13	40	67	12
Seldom found growing indoors**													
Ascospores	1,900	27	53	160	510	960	72	25	53	110	360	690	71
Basidiospores	800	53	110	430	1,800	3,900	95	53	80	260	990	2,300	93
Oidium	40	13	13	13	40	67	14	13	13	13	44	75	19
Rusts	-	8	13	13	40	73	14	13	13	13	53	80	26
Smuts, Periconia, Myxomycetes	-	13	13	27	67	110	54	13	13	40	110	210	68
§ TOTAL SPORES/m3	13,000												

[†]The "Typical Outdoor Data' represents the typical outdoor spore levels for the location and time frame indicated. The last column represents the frequency of occurrence. The very low, low, med, high, and very high values represent the 10, 20, 50, 80, and 90 percentile values of the spore type when it is detected. For example, if the frequency of occurrence is 63% and the low value is 53, it would mean that the given spore type is detected 63% of the time and, when detected, 20% of the time it is present in levels above the detection limit and below 53 spores/m3. These values are updated periodically, and if enough data is not available to make a statistically meaningful assessment, it is indicated with a dash.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor data" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. In addition, EMLab P&K may not have received and tested a representative number of samples for every region or time period. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the use or interpretation of the data contained in, or any actions taken or omitted in reliance upon, this report.

EMLab P&K, LLC EMLab ID: 1172421, Page 1 of 1

[§] Total Spores/m3 has been rounded to two significant figures to reflect analytical precision.

^{*} The spores in this category are generally capable of growing on wet building materials in addition to growing outdoors. Building related growth is dependent upon the fungal type, moisture level, type of material, and other factors. *Cladosporium* is one of the predominant spore types worldwide and is frequently present in high numbers. *Penicillium/Aspergillus* species colonize both outdoor and indoor wet surfaces rapidly and are very easily dispersed. Other genera are usually present in lesser numbers.

^{**} These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens. However, in each group there are notable exceptions. For example, agents of wood decay are members of the basidiomycetes and high counts of a single morphological type of basidiospore on an inside sample should be considered significant.

 $[\]ddagger$ n = number of samples used to calculate data.

Date of Sampling: 02-14-2014

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Date of Receipt: 02-18-2014 Re: 21402001-4 Date of Report: 02-19-2014

MoldSTATTM: Supplementary Statistical Spore Trap Report

Outdoor Summary: 21402001-4TM07OUT:

Species detected		Outdoor	r sample s	pores/m3	Typical outdoor ranges	Freq.
	<100	1K	10K	>100K	(North America)	%
Alternaria				27	7 - 33 - 590	45
Ascospores				1,900] 13 - 210 - 5,700	76
Basidiospores				800	17 - 450 - 24,000	92
Cladosporium				9,200	27 - 470 - 10,000	90
Epicoccum				40	7 - 20 - 340	25
Oidium				40	7 - 13 - 230	12
Penicillium/Aspergillus types				480	13 - 170 - 2,700	68
Smuts, Periconia, Myxomycetes				< 13	7 - 53 - 930	64
Stemphylium				67	7 - 13 - 93	3
Total				13,000		

The "Typical outdoor ranges" and "Freq. %" columns show the typical low, medium, and high spore counts per cubic meter and the frequency of occurrence for the given spore type. The low, medium, and high values represent the 2.5, 50, and 97.5 percentile values when the spore type is detected. For example, if the low value is 53 and the frequency of occurence is 63%, it would mean that we typically detect the given spore type on 63 percent of all outdoor samples and, when detected, 2.5% of the time it is present in levels below 53 spores/m3.

Indoor Samples

Location: 21402001-4TM08

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)		ent ratio** /outdoor)	correla	nan rank ation*** /outdoor)	MoldSCORE**** (indoor/outdoor)		
Result: < 1%	dF: 5 Result: 3.9524 Critical value: 11.0705 Inside Similar: Yes	Resul	t: 0.0000	Result Critical v	7: 10 : -0.2242 alue: 0.5515 Similar: No	Score: 113 Result: Lov		
Species 1	Detected			Spor	res/m3			
		<100	1K		10K	>100K		
	Other brown						27	
Smuts, Periconia, Myxomycetes							13	
	Total						40	

EMLab P&K, LLC EMLab ID: 1172421, Page 1 of 4

Date of Sampling: 02-14-2014

Client: Hygiene Technologies International, Inc. C/O: Mr. Lakhpreet Sandhu

Date of Receipt: 02-18-2014 Date of Report: 02-19-2014 Re: 21402001-4

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 21402001-4TM09

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman ra correlation* (indoor/outde	***	MoldSCORE**** (indoor/outdoor)		
Result: < 1%	dF: 5 Result: 3.9524 Critical value: 11.0705 Inside Similar: Yes	Result: 0.2222		dF: 8 Result: 0.6726 Critical value: 0.6190 Outside Similar: Yes		Score: 101 Result: Low		
Species 1	Detected	Spores/m3						
		<100	1K	10)K	>100K		
	Cladosporium					53		
	Total					53		

Location: 21402001-4TM10

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)		
Result: < 1%	dF: 5 Result: 3.9524 Critical value: 11.0705 Inside Similar: Yes	Result: 0.3636		dF: 9 Result: 0.3375 Critical value: 0.5833 Outside Similar: No	Score: 108 Result: Low		
Species 1	Detected			Spores/m3			
		<100	1K	10K	>100K		
	Cladosporium				53		
Other brown					13		
Penicillium/Aspergillus types					53		
	Total				120		

Location: 21402001-4TM11

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)			
Result: < 1%	dF: 5 Result: 3.9524 Critical value: 11.0705 Inside Similar: Yes	Result: 0.0000	dF: N/A Result: N/A Critical value: N/A Outside Similar: N/A	Score: 100 Result: Low			
Species 1	Detected	Spores/m3					
		<100 1K	10K	>100K			
	None Detected			< 13			

EMLab P&K, LLC EMLab ID: 1172421, Page 2 of 4

Date of Sampling: 02-14-2014

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Date of Receipt: 02-18-2014 Re: 21402001-4 Date of Report: 02-19-2014

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 21402001-4TM12

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)		
Result: 1%	dF: 5 Result: 3.9524 Critical value: 11.0705 Inside Similar: Yes	Result:	0.5455	dF: 8 Result: 0.8631 Critical value: 0.6190 Outside Similar: Yes	Score: 105 Result: Low		
Species 1	Detected			Spores/m3			
		<100	1K	10K	>100K		
	Ascospores				53		
	Basidiospores				53		
	Cladosporium				53		
	Total				160		

Location: 21402001-4TM13

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: < 1%	dF: 5 Result: 3.9524 Critical value: 11.0705 Inside Similar: Yes	Result: 0.0000	dF: N/A Result: N/A Critical value: N/A Outside Similar: N/A	Score: 100 Result: Low
Species	Detected		Spores/m3	
		<100 1K	10K	>100K
	None Detected			< 13

^{*} The Friedman chi-square statistic is a non-parametric test that examines variation in a set of data (in this case, all indoor spore counts). The null hypothesis (H0) being tested is that there is no meaningful difference in the data for all indoor locations. The alternative hypothesis (used if the test disproves the null hypothesis) is that there is a difference between the indoor locations. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

EMLab P&K, LLC EMLab ID: 1172421, Page 3 of 4

^{**} An agreement ratio is a simple method for assessing the similarity of two samples (in this case the indoor sample and the outdoor summary) based on the spore types present. A score of one indicates that the types detected in one location are the same as that in the other. A score of zero indicates that none of the types detected indoors are present outdoors. Typically, an agreement of 0.8 or higher is considered high.

^{***} The Spearman rank correlation is a non-parametric test that examines correlation between two sets of data (in this case the indoor location and the outdoor summary). The null hypothesis (H0) being tested is that the indoor and outdoor samples are unrelated. The alternative hypothesis (used if the test disproves the null hypothesis) is that the samples are similar. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-14-2014 Date of Receipt: 02-18-2014 Date of Report: 02-19-2014

MoldSTATTM: Supplementary Statistical Spore Trap Report

**** MoldSCORETM is a specialized method for examining air sampling data. It is a score between 100 and 300, with 100 indicating a greater likelihood that the airborne indoor spores originated from the outside, and 300 indicating a greater likelihood that they originated from an inside source. The Result displayed is based on the numeric score given and will be either Low, Medium, or High, indicating a low, medium, or high likelihood that the spores detected originated from an indoor source. EMLab P&Kreserves the right to, and may at anytime, modify or change the MoldScore algorithm without notice.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor ranges" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. With the statistical analysis provided, as with all statistical comparisons and analyses, false-positive and false-negative results can and do occur. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the data contained in, or any actions taken or omitted in reliance upon, this report.

EMLab P&K, LLC EMLab ID: 1172421, Page 4 of 4

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu Re: 21402001-4

Date of Sampling: 02-14-2014 Date of Receipt: 02-18-2014 Date of Report: 02-19-2014

MoldSCORETM: **Spore Trap Report Outdoor Sample:** 21402001-4TM07OUT

Fungi Identified	Ou	tdo	Raw	Spores/				
_	<100)	1K		10K	>1001	count	m3
Generally able to grow indoors*								
Alternaria							2	27
Bipolaris/Drechslera group							ND	< 13
Chaetomium				Ш			ND	< 13
Cladosporium				Ш			173	9,200
Curvularia							ND	< 13
Epicoccum				Ш			3	40
Nigrospora							ND	< 13
Penicillium/Aspergillus types†							9	480
Stachybotrys				Ш			ND	< 13
Stemphylium							5	67
Torula							ND	< 13
Seldom found growing indoors**								
Ascospores				Ш			36	1,900
Basidiospores							15	800
Oidium				Ш			3	40
Rusts				Ш			ND	< 13
Smuts, Periconia, Myxomycetes				Ш			ND	< 13
Total								12,600

Fungi Identified	In	door sa	mple :	spor	es/n	13	Raw	Spores/
	<100	1K		10K	>	100K	count	m3
Generally able to grow indoors*								
Alternaria							ND	< 13
Bipolaris/Drechslera group							ND	< 13
Chaetomium							ND	< 13
Cladosporium							ND	< 13
Curvularia							ND	< 13
Nigrospora							ND	< 13
Other brown							2	27
Penicillium/Aspergillus types†							ND	< 13
Stachybotrys							ND	< 13
Torula							ND	< 13
Seldom found growing indoors**								
Ascospores							ND	< 13
Basidiospores							ND	< 13
Rusts							ND	< 13
Smuts, Periconia, Myxomycetes							1	13
Total								40

100	MoldSCORE; 200 300 Score											
			100									
			100									
			100									
			100									
			100									
			100									
			111									
			100									
			100									
			100									
			100									
			100									
			100									
			103									
Fina	113											

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-14-2014 Date of Receipt: 02-18-2014 Date of Report: 02-19-2014

 $\mathbf{MoldSCORE}^{\mathsf{TM}}\mathbf{:}\ \mathbf{Spore}\ \mathbf{Trap}\ \mathbf{Report}$

Location: 21402001-4TM09

Fungi Identified	Ind	loo	r	sam	Raw	Spores/					
_	<100			1K		10K	>	-100	K	count	m3
Generally able to grow indoors*											
Alternaria										ND	< 13
Bipolaris/Drechslera group										ND	< 13
Chaetomium										ND	< 13
Cladosporium										1	53
Curvularia										ND	< 13
Nigrospora										ND	< 13
Penicillium/Aspergillus types†										ND	< 13
Stachybotrys										ND	< 13
Torula										ND	< 13
Seldom found growing indoors**											
Ascospores										ND	< 13
Basidiospores										ND	< 13
Rusts										ND	< 13
Smuts, Periconia, Myxomycetes										ND	< 13
Total				•							53

100	100 MoldSCORE : 300 300											
			100									
			100									
			100									
			101									
			100									
			100									
			100									
			100									
			100									
			100									
			100									
			100									
			100									
Fina	Final MoldSCORE											

Fungi Identified	In	Indoor sample spores/m3										Raw	Spores/
	<100)		1K			1	0K		>100	K	count	m3
Generally able to grow indoors*													
Alternaria												ND	< 13
Bipolaris/Drechslera group												ND	< 13
Chaetomium												ND	< 13
Cladosporium												1	53
Curvularia												ND	< 13
Nigrospora												ND	< 13
Other brown												1	13
Penicillium/Aspergillus types†												1	53
Stachybotrys												ND	< 13
Torula												ND	< 13
Seldom found growing indoors**													
Ascospores												ND	< 13
Basidiospores												ND	< 13
Rusts												ND	< 13
Smuts, Periconia, Myxomycetes												ND	< 13
Total													120

100	100 MoldSCORE; 300 Score												
			100										
			100										
			100										
			100										
			100										
			100										
			105										
			108										
			100										
			100										
			100										
			100										
			100										
			100										
Fina	al MoldSCO	RE	108										

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-14-2014 Date of Receipt: 02-18-2014 Date of Report: 02-19-2014

 $\mathbf{MoldSCORE}^{\mathsf{TM}}\mathbf{:}\ \mathbf{Spore}\ \mathbf{Trap}\ \mathbf{Report}$

Location: 21402001-4TM11

Fungi Identified	Ind	00	r sa	mp	Raw	Spores/			
_	<100		1 K	: _	10K	>	1001	count	m3
Generally able to grow indoors*									
Alternaria								ND	< 13
Bipolaris/Drechslera group								ND	< 13
Chaetomium								ND	< 13
Cladosporium								ND	< 13
Curvularia								ND	< 13
Nigrospora								ND	< 13
Penicillium/Aspergillus types†								ND	< 13
Stachybotrys								ND	< 13
Torula		$\ $						ND	< 13
Seldom found growing indoors**									
Ascospores								ND	< 13
Basidiospores								ND	< 13
Rusts								ND	< 13
Smuts, Periconia, Myxomycetes								ND	< 13
Total									N/A

100	100 MoldSCORE : 300 300										
	_										
			100								
			100								
			100								
			100								
			100								
			100								
			100								
			100								
			100								
			100								
			100								
			100								
			100								
Fina	Final MoldSCORE										

Fungi Identified	In	do	or	san	Raw	Spores/				
	<100)		1K		10K	>10)0K	count	m3
Generally able to grow indoors*										
Alternaria									ND	< 13
Bipolaris/Drechslera group									ND	< 13
Chaetomium								Ш	ND	< 13
Cladosporium									1	53
Curvularia									ND	< 13
Nigrospora									ND	< 13
Penicillium/Aspergillus types†									ND	< 13
Stachybotrys									ND	< 13
Torula									ND	< 13
Seldom found growing indoors**										
Ascospores									1	53
Basidiospores									1	53
Rusts									ND	< 13
Smuts, Periconia, Myxomycetes									ND	< 13
Total										160

_																	
1	MoldSCORE 200 300																
Г																	
																	100
																	100
	1																100
																	100
																	100
																	100
																	100
																	100
																	100
L					_												
																	112
																	105
																	100
																	100
]	Final MoldSCORE									105							

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-14-2014 Date of Receipt: 02-18-2014 Date of Report: 02-19-2014

MoldSCORETM: Spore Trap Report

Location: 21402001-4TM13

Fungi Identified	Ind	Indoor sample spores/m3							Raw	Spores/
_	<100		11	K		10K	>	1001	count	m3
Generally able to grow indoors*										
Alternaria									ND	< 13
Bipolaris/Drechslera group									ND	< 13
Chaetomium									ND	< 13
Cladosporium									ND	< 13
Curvularia									ND	< 13
Nigrospora									ND	< 13
Penicillium/Aspergillus types†									ND	< 13
Stachybotrys									ND	< 13
Torula									ND	< 13
Seldom found growing indoors**										
Ascospores									ND	< 13
Basidiospores									ND	< 13
Rusts									ND	< 13
Smuts, Periconia, Myxomycetes					Ш				ND	< 13
Total										N/A

l N										
100	100 200 300									
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
Final	MoldSC	ORE	100							

^{*} The spores in this category are generally capable of growing on wet building materials in addition to growing outdoors. Building related growth is dependent upon the fungal type, moisture level, type of material, and other factors. *Cladosporium* is one of the predominant spore types worldwide and is frequently present in high numbers. *Penicillium/Aspergillus* species colonize both outdoor and indoor wet surfaces rapidly and are very easily dispersed. Other genera are usually present in lesser numbers.

EMLab P&K, LLC EMLab ID: 1172421, Page 4 of 4

^{**} These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens. However, in each group there are notable exceptions. For example, agents of wood decay are members of the basidiomycetes and high counts of a single morphological type of basidiospore on an inside sample should be considered significant.

[†]The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods.

[‡]Rated on a scale from 100 to 300. A rating less than 150 is low and indicates a low probability of spores originating inside. A rating greater than 250 is high and indicates a high probability that the spores originated from inside, presumably from indoor mold growth. A rating between 150 and 250 indicates a moderate likelihood of indoor fungal growth. MoldSCORE is NOT intended for wall cavity samples. It is intended for ambient air samples in residences. Using the analysis on other samples (like wall cavity samples) will lead to misleading results.



Report for:

Mr. Kenny Hsi, Mr. Lakhpreet Sandhu Hygiene Technologies International, Inc. 3625 Del Amo Boulevard, Suite 180 Torrance, CA 90503-8370

Regarding: Project: 21402001-4

EMĹ ID: 1173747

Approved by:

Technical Manager Melissa Tracey

Service SOPs: Spore trap analysis (1038) AIHA-LAP, LLC accredited service, Lab ID #179768

All samples were received in acceptable condition unless noted in the Report Comments portion in the body of the report. Due to the nature of the analyses performed, field blank correction of results is not applied. The results relate only to the items tested.

Dates of Analysis:

Spore trap analysis: 02-21-2014

EMLab P&K ("the Company") shall have no liability to the client or the client's customer with respect to decisions or recommendations made, actions taken or courses of conduct implemented by either the client or the client's customer as a result of or based upon the Test Results. In no event shall the Company be liable to the client with respect to the Test Results except for the Company's own willful misconduct or gross negligence nor shall the Company be liable for incidental or consequential damages or lost profits or revenues to the fullest extent such liability may be disclaimed by law, even if the Company has been advised of the possibility of such damages, lost profits or lost revenues. In no event shall the Company's liability with respect to the Test Results exceed the amount paid to the Company by the client therefor.

Client: Hygiene Technologies International, Inc. Date of Sampling: 02-20-2014 C/O: Mr. Kenny Hsi, Mr. Lakhpreet Sandhu Re: 21402001-4

Date of Receipt: 02-21-2014 Date of Report: 02-21-2014

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:	21402001	-4TM14OUT	214020	01-4TM15	21402001-4TM16		
Comments (see below)	None		N	Vone	None		
Lab ID-Version‡:	5314431-1		531	4432-1	5314433-1		
Analysis Date:	02/21/2014		02/2	21/2014	02/21/2014		
	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3	
Alternaria	5	67		_		_	
Ascospores	3	160					
Basidiospores	19	1,000	4	210	1	53	
Botrytis		·					
Chaetomium							
Cladosporium	81	4,300	6	320	4	210	
Curvularia	1	13					
Epicoccum	1	13					
Fusarium							
Myrothecium							
Nigrospora							
Other colorless							
Penicillium/Aspergillus types†	6	320					
Pithomyces	1	13					
Rusts	1	13					
Smuts, Periconia, Myxomycetes	3	40	1	13			
Stachybotrys							
Stemphylium							
Torula							
Ulocladium							
Zygomycetes							
Background debris (1-4+)††	2+		2+		2+		
Hyphal fragments/m3	93		40		< 13		
Pollen/m3	160		13		< 13		
Skin cells (1-4+)	< 1+		1+		1+		
Sample volume (liters)	75		75		75		
§ TOTAL SPORES/m3		6,000		550		270	

Comments:

Spore types listed without a count or data entry were not detected during the course of the analysis for the respective sample.

The analytical sensitivity is the spores/m3 divided by the raw count. The limit of detection is the analytical sensitivity multiplied by the sample volume divided by 1000.

EMLab P&K, LLC EMLab ID: 1173747, Page 2 of 3

[†] The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††}Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be higher then reported. It is important to account for samples volumes when evaluating dust levels.

For more information regarding analytical sensitivity, please contact QA by calling the laboratory. ‡ A "Version" indicated by -"x" after the Lab ID# with a value greater than 1 indicates a sample with amended data. The revision number is reflected by the value of "x".

[§] Total Spores/m3 has been rounded to two significant figures to reflect analytical precision.

Client: Hygiene Technologies International, Inc. C/O: Mr. Kenny Hsi, Mr. Lakhpreet Sandhu Re: 21402001-4

Date of Sampling: 02-20-2014 Date of Receipt: 02-21-2014 Date of Report: 02-21-2014

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:	214020	01-4TM17	214020	01-4TM18	21402001-4TM19		
Comments (see below)	None		N	Vone	None		
Lab ID-Version‡:	5314434-1		531	4435-1	5314436-1		
Analysis Date:	02/21/2014		02/2	21/2014	02/21/2014		
	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3	
Alternaria		_	1	13		_	
Ascospores							
Basidiospores	1	53	4	210	1	53	
Botrytis							
Chaetomium							
Cladosporium	1	53	7	370	1	53	
Curvularia							
Epicoccum							
Fusarium							
Myrothecium							
Nigrospora							
Other colorless							
Penicillium/Aspergillus types†					2	110	
Pithomyces							
Rusts							
Smuts, Periconia, Myxomycetes							
Stachybotrys							
Stemphylium							
Torula							
Ulocladium							
Zygomycetes							
Background debris (1-4+)††	2+		2+		2+		
Hyphal fragments/m3	13		27		< 13		
Pollen/m3	< 13		< 13		< 13		
Skin cells (1-4+)	1+		1+		1+		
Sample volume (liters)	75		75		75		
§ TOTAL SPORES/m3		110		600		210	

Comments:

Spore types listed without a count or data entry were not detected during the course of the analysis for the respective sample.

The analytical sensitivity is the spores/m3 divided by the raw count. The limit of detection is the analytical sensitivity multiplied by the sample volume divided by 1000.

EMLab P&K, LLC EMLab ID: 1173747, Page 3 of 3

[†] The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††}Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be higher then reported. It is important to account for samples volumes when evaluating dust levels.

For more information regarding analytical sensitivity, please contact QA by calling the laboratory. ‡ A "Version" indicated by -"x" after the Lab ID# with a value greater than 1 indicates a sample with amended data. The revision number is reflected by the value of "x".

[§] Total Spores/m3 has been rounded to two significant figures to reflect analytical precision.

Client: Hygiene Technologies International, Inc. C/O: Mr. Kenny Hsi, Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-20-2014 Date of Receipt: 02-21-2014 Date of Report: 02-21-2014

$MoldRANGE^{TM}\hbox{:} \ Extended \ Outdoor \ Comparison$

Outdoor Location: 21402001-4TM14OUT

Fungi Identified	Outdoor	Typical Outdoor Data for:					Typical Outdoor Data for:						
	data	February in California† (n‡=16302)					The entire year in California† (n‡=200710)						
	spores/m3	very low	low	med	high	very high	freq %	very low	low	med	high	very high	freq %
Generally able to grow indoors*													
Alternaria	67	13	13	20	44	67	38	13	13	27	67	110	53
Bipolaris/Drechslera group	-	7	13	13	27	40	7	7	13	13	27	40	12
Chaetomium	-	7	13	13	27	40	9	8	13	13	27	47	19
Cladosporium	4,300	80	150	400	1,000	1,700	95	110	210	610	1,600	2,800	97
Curvularia	13	7	10	13	13	27	2	7	13	13	27	53	6
Epicoccum	13	7	13	13	27	53	13	8	13	13	33	53	19
Nigrospora	-	7	13	13	13	27	4	7	13	13	27	53	8
Penicillium/Aspergillus types	320	53	66	200	490	820	82	53	100	210	590	1,000	84
Pithomyces	13	7	10	13	13	35	1	7	13	13	27	53	4
Stachybotrys	-	13	13	13	40	80	3	7	13	13	33	67	4
Torula	-	7	13	13	40	53	5	8	13	13	40	67	12
Seldom found growing indoors**													
Ascospores	160	27	53	160	510	960	72	25	53	110	360	690	71
Basidiospores	1,000	53	110	430	1,800	3,900	95	53	80	260	990	2,300	93
Rusts	13	8	13	13	40	73	14	13	13	13	53	80	26
Smuts, Periconia, Myxomycetes	40	13	13	27	67	110	54	13	13	40	110	210	68
§ TOTAL SPORES/m3	6,000												

†The 'Typical Outdoor Data' represents the typical outdoor spore levels for the location and time frame indicated. The last column represents the frequency of occurrence. The very low, low, med, high, and very high values represent the 10, 20, 50, 80, and 90 percentile values of the spore type when it is detected. For example, if the frequency of occurrence is 63% and the low value is 53, it would mean that the given spore type is detected 63% of the time and, when detected, 20% of the time it is present in levels above the detection limit and below 53 spores/m3. These values are updated periodically, and if enough data is not available to make a statistically meaningful assessment, it is indicated with a dash

§ Total Spores/m3 has been rounded to two significant figures to reflect analytical precision.

 \ddagger n = number of samples used to calculate data.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor data" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. In addition, EMLab P&K may not have received and tested a representative number of samples for every region or time period. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the use or interpretation of the data contained in, or any actions taken or omitted in reliance upon, this report.

EMLab P&K, LLC EMLab ID: 1173747, Page 1 of 1

^{*} The spores in this category are generally capable of growing on wet building materials in addition to growing outdoors. Building related growth is dependent upon the fungal type, moisture level, type of material, and other factors. *Cladosporium* is one of the predominant spore types worldwide and is frequently present in high numbers. *Penicillium/Aspergillus* species colonize both outdoor and indoor wet surfaces rapidly and are very easily dispersed. Other genera are usually present in lesser numbers.

^{**} These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens. However, in each group there are notable exceptions. For example, agents of wood decay are members of the basidiomycetes and high counts of a single morphological type of basidiospore on an inside sample should be considered significant.

Client: Hygiene Technologies International, Inc. C/O: Mr. Kenny Hsi, Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-20-2014 Date of Receipt: 02-21-2014 Date of Report: 02-21-2014

MoldSTATTM: Supplementary Statistical Spore Trap Report

Outdoor Summary: 21402001-4TM14OUT:

Species detected		Outdoo	r sample s	pores/m3	1	ГурісаІ	outdo	or ranges	Freq.
	<100	1K	10K	>100K		(Nor	th An	nerica)	%
Alternaria				67		7 -	33	- 590	45
Ascospores				160		13 -	210	- 5,700	76
Basidiospores				1,000		17 -	450	- 24,000	92
Cladosporium				4,300		27 -	470	- 10,000	90
Curvularia				13		7 -	27	- 600	17
Epicoccum				13		7 -	20	- 340	25
Penicillium/Aspergillus types				320		13 -	170	- 2,700	68
Pithomyces				13		7 -	20	- 590	15
Rusts				13		7 -	20	- 360	20
Smuts, Periconia, Myxomycetes				40		7 -	53	- 930	64
Total				6,000					

The "Typical outdoor ranges" and "Freq. %" columns show the typical low, medium, and high spore counts per cubic meter and the frequency of occurrence for the given spore type. The low, medium, and high values represent the 2.5, 50, and 97.5 percentile values when the spore type is detected. For example, if the low value is 53 and the frequency of occurrence is 63%, it would mean that we typically detect the given spore type on 63 percent of all outdoor samples and, when detected, 2.5% of the time it is present in levels below 53 spores/m3.

Indoor Samples

Location: 21402001-4TM15

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)						
Result: 9%	dF: 4 Result: 4.5600 Critical value: 9.4877 Inside Similar: Yes	Result: 0.4615	dF: 10 Result: 0.7152 Critical value: 0.5515 Outside Similar: Yes	Score: 113 Result: Low						
Species	Detected	Spores/m3								
		<100 1K	10K	>100K						
	Basidiospores			210						
	Cladosporium			320						
Smuts, Periconia, Myxomycetes				13						
	Total			550						

EMLab P&K, LLC EMLab ID: 1173747, Page 1 of 3

Client: Hygiene Technologies International, Inc. C/O: Mr. Kenny Hsi, Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-20-2014 Date of Receipt: 02-21-2014 Date of Report: 02-21-2014

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 21402001-4TM16

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)		
Result: 4%	dF: 4 Result: 4.5600 Critical value: 9.4877 Inside Similar: Yes	Result: 0.3333	dF: 10 Result: 0.7758 Critical value: 0.5515 Outside Similar: Yes	Score: 101 Result: Low		
Species	Detected		Spores/m3			
		<100 1K	10K	>100K		
	Basidiospores			53		
	Cladosporium			210		
	Total			270		

Location: 21402001-4TM17

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)		
Result: 1%	dF: 4 Result: 4.5600 Critical value: 9.4877 Inside Similar: Yes	Result: 0.3333	dF: 10 Result: 0.7727 Critical value: 0.5515 Outside Similar: Yes	Score: 104 Result: Low		
Species 1	Detected		Spores/m3			
_		<100 1K	10K	>100K		
	Basidiospores			53		
	Cladosporium			53		
	Total			110		

Location: 21402001-4TM18

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)						
Result: 9%	dF: 4 Result: 4.5600 Critical value: 9.4877 Inside Similar: Yes	Result: 0.4615	dF: 10 Result: 0.7636 Critical value: 0.5515 Outside Similar: Yes	Score: 112 Result: Low						
Species 1	Detected	Spores/m3								
		<100 1K	10K	>100K						
	Alternaria			13						
	Basidiospores			210						
	Cladosporium			370						
	Total			600						

EMLab ID: 1173747, Page 2 of 3

Client: Hygiene Technologies International, Inc. C/O: Mr. Kenny Hsi, Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-20-2014 Date of Receipt: 02-21-2014 Date of Report: 02-21-2014

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 21402001-4TM19

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)		nt ratio** outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)			
Result: 3%	dF: 4 Result: 4.5600 Critical value: 9.4877 Inside Similar: Yes	Result:	0.4615	dF: 10 Result: 0.8212 Critical value: 0.5515 Outside Similar: Yes	Score: 116 Result: Low			
Species	Detected			Spores/m3				
		<100	1K	10K	>100K			
	Basidiospores				53			
	Cladosporium				53			
Penicillium/Aspergillus types					110			
	Total				210			

^{*} The Friedman chi-square statistic is a non-parametric test that examines variation in a set of data (in this case, all indoor spore counts). The null hypothesis (H0) being tested is that there is no meaningful difference in the data for all indoor locations. The alternative hypothesis (used if the test disproves the null hypothesis) is that there is a difference between the indoor locations. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

**** MoldSCORETM is a specialized method for examining air sampling data. It is a score between 100 and 300, with 100 indicating a greater likelihood that the airborne indoor spores originated from the outside, and 300 indicating a greater likelihood that they originated from an inside source. The Result displayed is based on the numeric score given and will be either Low, Medium, or High, indicating a low, medium, or high likelihood that the spores detected originated from an indoor source. EMLab P&Kreserves the right to, and may at anytime, modify or change the MoldScore algorithm without notice.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor ranges" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. With the statistical analysis provided, as with all statistical comparisons and analyses, false-positive and false-negative results can and do occur. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the data contained in, or any actions taken or omitted in reliance upon, this report.

EMLab P&K, LLC EMLab ID: 1173747, Page 3 of 3

^{**} An agreement ratio is a simple method for assessing the similarity of two samples (in this case the indoor sample and the outdoor summary) based on the spore types present. A score of one indicates that the types detected in one location are the same as that in the other. A score of zero indicates that none of the types detected indoors are present outdoors. Typically, an agreement of 0.8 or higher is considered high.

^{***} The Spearman rank correlation is a non-parametric test that examines correlation between two sets of data (in this case the indoor location and the outdoor summary). The null hypothesis (H0) being tested is that the indoor and outdoor samples are unrelated. The alternative hypothesis (used if the test disproves the null hypothesis) is that the samples are similar. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

Client: Hygiene Technologies International, Inc. C/O: Mr. Kenny Hsi, Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-20-2014 Date of Receipt: 02-21-2014 Date of Report: 02-21-2014

MoldSCORETM: **Spore Trap Report Outdoor Sample:** 21402001-4TM14OUT

Fungi Identified	Οι	ıtde	or	sam	ıple	spo	res	/m3	Raw	Spores/
	<10	0		١K		10K		>100 F	count	m3
Generally able to grow indoors*										
Alternaria									5	67
Bipolaris/Drechslera group									ND	< 13
Chaetomium									ND	< 13
Cladosporium									81	4,300
Curvularia									1	13
Epicoccum									1	13
Nigrospora									ND	< 13
Penicillium/Aspergillus types†									6	320
Pithomyces									1	13
Stachybotrys									ND	< 13
Torula									ND	< 13
Seldom found growing indoors**										
Ascospores									3	160
Basidiospores									19	1,000
Rusts									1	13
Smuts, Periconia, Myxomycetes									3	40
Total										5,973

Fungi Identified	In	do	or	sam	ple	S	por	es/ı	m3		Raw	Spores/
	<10	0		1K			10K	>	>100	K	count	m3
Generally able to grow indoors*												
Alternaria											ND	< 13
Bipolaris/Drechslera group											ND	< 13
Chaetomium											ND	< 13
Cladosporium											6	320
Curvularia											ND	< 13
Nigrospora											ND	< 13
Penicillium/Aspergillus types†											ND	< 13
Stachybotrys											ND	< 13
Torula											ND	< 13
Seldom found growing indoors**												
Ascospores											ND	< 13
Basidiospores											4	210
Rusts											ND	< 13
Smuts, Periconia, Myxomycetes											1	13
Total												547

100	MoldSCORE 200 300									
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			113							
			100							
			102							
Fina	Final MoldSCORE									

Client: Hygiene Technologies International, Inc. C/O: Mr. Kenny Hsi, Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-20-2014 Date of Receipt: 02-21-2014 Date of Report: 02-21-2014

MoldSCORETM: Spore Trap Report

Location: 21402001-4TM16

Fungi Identified	Inde	o	sam	ıple	sp	ore	s/n	13	Raw	Spores/
	<100		1K		1	0K	>	100I	count	m3
Generally able to grow indoors*										
Alternaria									ND	< 13
Bipolaris/Drechslera group									ND	< 13
Chaetomium									ND	< 13
Cladosporium									4	210
Curvularia									ND	< 13
Nigrospora					\prod				ND	< 13
Penicillium/Aspergillus types†									ND	< 13
Stachybotrys									ND	< 13
Torula		Ш			\prod				ND	< 13
Seldom found growing indoors**										
Ascospores		Ш			Ш				ND	< 13
Basidiospores									1	53
Rusts		П							ND	< 13
Smuts, Periconia, Myxomycetes									ND	< 13
Total										267

100	MoldSC 200	ORE:	
			100
			100
			100
			101
			100
			100
			100
			100
			100
			100
			101
			100
			100
Fina	d MoldSC	ORE	101

Fungi Identified	In	do	or	S	amj	ple	S	por	es/	/m	3	Raw	Spores/
	<10	C		1	K			10K		>1	00K	count	m3
Generally able to grow indoors*													
Alternaria												ND	< 13
Bipolaris/Drechslera group			Ш									ND	< 13
Chaetomium												ND	< 13
Cladosporium												1	53
Curvularia												ND	< 13
Nigrospora												ND	< 13
Penicillium/Aspergillus types†												ND	< 13
Stachybotrys												ND	< 13
Torula												ND	< 13
Seldom found growing indoors**													
Ascospores												ND	< 13
Basidiospores												1	53
Rusts												ND	< 13
Smuts, Periconia, Myxomycetes												ND	< 13
Total													107

100	MoldSCORE‡ 200 300 Score								
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			104						
			100						
			100						
Fina	Final MoldSCORE								

Client: Hygiene Technologies International, Inc. C/O: Mr. Kenny Hsi, Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-20-2014 Date of Receipt: 02-21-2014 Date of Report: 02-21-2014

MoldSCORETM: Spore Trap Report

Location: 21402001-4TM18

Fungi Identified	Inc	doc	or	sam	ple	S	pore	s/ı	m3	}	Raw	Spores/
	<100			1K			10K	>	>100)K	count	m3
Generally able to grow indoors*												
Alternaria											1	13
Bipolaris/Drechslera group											ND	< 13
Chaetomium											ND	< 13
Cladosporium											7	370
Curvularia											ND	< 13
Nigrospora											ND	< 13
Penicillium/Aspergillus types†											ND	< 13
Stachybotrys											ND	< 13
Torula											ND	< 13
Seldom found growing indoors**												
Ascospores											ND	< 13
Basidiospores											4	210
Rusts											ND	< 13
Smuts, Periconia, Myxomycetes											ND	< 13
Total				•			•					600

MoldSCORE	
100 200 300	Score
	102
	100
	100
	100
	100
	100
	100
	100
	100
	100
	112
	100
	100
Final MoldSCORE	112

Fungi Identified	ln	Indoor sample spores/m3							3	Raw	Spores/	
	<100)	1	K			10K		>10	00K	count	m3
Generally able to grow indoors*												
Alternaria											ND	< 13
Bipolaris/Drechslera group											ND	< 13
Chaetomium											ND	< 13
Cladosporium											1	53
Curvularia											ND	< 13
Nigrospora											ND	< 13
Penicillium/Aspergillus types†											2	110
Stachybotrys											ND	< 13
Torula											ND	< 13
Seldom found growing indoors**												
Ascospores											ND	< 13
Basidiospores											1	53
Rusts											ND	< 13
Smuts, Periconia, Myxomycetes											ND	< 13
Total												213

100	MoldSCORE 100 200 30								
100	200	500	Беоге						
			100						
			100						
			100						
			100						
			100						
			100						
			116						
			100						
			100						
			100						
			102						
			100						
			100						
Fina	Final MoldSCORE								

Client: Hygiene Technologies International, Inc. C/O: Mr. Kenny Hsi, Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-20-2014 Date of Receipt: 02-21-2014 Date of Report: 02-21-2014

MoldSCORETM: Spore Trap Report

* The spores in this category are generally capable of growing on wet building materials in addition to growing outdoors. Building related growth is dependent upon the fungal type, moisture level, type of material, and other factors. *Cladosporium* is one of the predominant spore types worldwide and is frequently present in high numbers. *Penicillium/Aspergillus* species colonize both outdoor and indoor wet surfaces rapidly and are very easily dispersed. Other genera are usually present in lesser numbers.

** These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens. However, in each group there are notable exceptions. For example, agents of wood decay are members of the basidiomycetes and high counts of

a single morphological type of basidiospore on an inside sample should be considered significant.

†The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods.

‡Rated on a scale from 100 to 300. A rating less than 150 is low and indicates a low probability of spores originating inside. A rating greater than 250 is high and indicates a high probability that the spores originated from inside, presumably from indoor mold growth. A rating between 150 and 250 indicates a moderate likelihood of indoor fungal growth. MoldSCORE is NOT intended for wall cavity samples. It is intended for ambient air samples in residences. Using the analysis on other samples (like wall cavity samples) will lead to misleading results.

EMLab P&K, LLC EMLab ID: 1173747, Page 4 of 4



Report for:

Mr. Lakhpreet Sandhu Hygiene Technologies International, Inc. 3625 Del Amo Boulevard, Suite 180 Torrance, CA 90503-8370

Regarding: Project: 21402001-4

EMĹ ID: 1175106

Approved by:

' ()

Technical Manager Melissa Tracey Dates of Analysis:

Spore trap analysis: 02-26-2014

Service SOPs: Spore trap analysis (1038) AIHA-LAP, LLC accredited service, Lab ID #179768

All samples were received in acceptable condition unless noted in the Report Comments portion in the body of the report. Due to the nature of the analyses performed, field blank correction of results is not applied. The results relate only to the items tested.

EMLab P&K ("the Company") shall have no liability to the client or the client's customer with respect to decisions or recommendations made, actions taken or courses of conduct implemented by either the client or the client's customer as a result of or based upon the Test Results. In no event shall the Company be liable to the client with respect to the Test Results except for the Company's own willful misconduct or gross negligence nor shall the Company be liable for incidental or consequential damages or lost profits or revenues to the fullest extent such liability may be disclaimed by law, even if the Company has been advised of the possibility of such damages, lost profits or lost revenues. In no event shall the Company's liability with respect to the Test Results exceed the amount paid to the Company by the client therefor.

Date of Sampling: 02-24-2014

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Date of Receipt: 02-25-2014 Re: 21402001-4 Date of Report: 02-26-2014

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:		02001- 20OUT	214020	01-4TM21	214020	01-4TM22	214020	01-4TM23	
Comments (see below)	N	Vone	N	lone	N	Vone	None		
Lab ID-Version‡:	532	0638-1	532	0639-1	532	0640-1	5320641-1		
Analysis Date:	02/26/2014		02/2	6/2014	02/2	26/2014	02/2	6/2014	
	raw ct.	spores/m3	raw ct.	spores/m3	raw ct. spores/m3		raw ct.	spores/m3	
Alternaria	3	40		•				•	
Ascospores	11	590							
Basidiospores	24	1,300					4	210	
Chaetomium		,							
Cladosporium	90	4,800					2	110	
Epicoccum									
Fusarium									
Myrothecium									
Nigrospora									
Oidium									
Other brown	1	13					1	13	
Other colorless									
Penicillium/Aspergillus types†	2	110							
Pithomyces									
Rusts	4	53							
Smuts, Periconia, Myxomycetes	6	80							
Stachybotrys									
Stemphylium	1	13							
Torula	6	80							
Ulocladium									
Zygomycetes									
Background debris (1-4+)††	2+		2+		2+		2+		
Hyphal fragments/m3	53		< 13		< 13		< 13		
Pollen/m3	67		13		< 13		27		
Skin cells (1-4+)	< 1+		1+		1+		1+		
Sample volume (liters)	75		75		75		75		
§ TOTAL SPORES/m3		7,100		< 13		< 13		330	

Comments:

Spore types listed without a count or data entry were not detected during the course of the analysis for the respective sample.

The analytical sensitivity is the spores/m3 divided by the raw count. The limit of detection is the analytical sensitivity multiplied by the sample volume divided by 1000.

EMLab P&K, LLC EMLab ID: 1175106, Page 2 of 3

[†] The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††}Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be higher then reported. It is important to account for samples volumes when evaluating dust levels.

For more information regarding analytical sensitivity, please contact QA by calling the laboratory. \ddagger A "Version" indicated by -"x" after the Lab ID# with a value greater than 1 indicates a sample with amended data. The revision number is reflected by the value of "x".

[§] Total Spores/m3 has been rounded to two significant figures to reflect analytical precision.

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu Re: 21402001-4

Date of Sampling: 02-24-2014 Date of Receipt: 02-25-2014 Date of Report: 02-26-2014

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:	214020	01-4TM24	214020	01-4TM25	214020	01-4TM26
Comments (see below)	ľ	None	N	Vone	1	None
Lab ID-Version‡:	532	0642-1	532	0643-1	532	20644-1
Analysis Date:	02/2	26/2014	02/2	26/2014	02/2	26/2014
	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3
Alternaria		_		_		_
Ascospores						
Basidiospores			1	53		
Chaetomium						
Cladosporium	1	53	2	110	1	53
Curvularia						
Epicoccum						
Fusarium						
Myrothecium						
Nigrospora						
Oidium	1	13				
Other brown	1	13				
Other colorless						
Penicillium/Aspergillus types†					1	53
Pithomyces						
Rusts	1	13	1	13		
Smuts, Periconia, Myxomycetes						
Stachybotrys						
Stemphylium						
Torula						
Ulocladium						
Zygomycetes						
Background debris (1-4+)††	2+		2+		2+	
Hyphal fragments/m3	< 13		< 13		< 13	
Pollen/m3	< 13		< 13		< 13	
Skin cells (1-4+)	1+		1+		1+	
Sample volume (liters)	75		75		75	
§ TOTAL SPORES/m3		93		170		110

Comments:

Spore types listed without a count or data entry were not detected during the course of the analysis for the respective sample.

The analytical sensitivity is the spores/m3 divided by the raw count. The limit of detection is the analytical sensitivity multiplied by the sample volume divided by 1000.

EMLab P&K, LLC EMLab ID: 1175106, Page 3 of 3

[†] The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††}Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be higher then reported. It is important to account for samples volumes when evaluating dust levels.

For more information regarding analytical sensitivity, please contact QA by calling the laboratory. ‡ A "Version" indicated by -"x" after the Lab ID# with a value greater than 1 indicates a sample with amended data. The revision number is reflected by the value of "x".

[§] Total Spores/m3 has been rounded to two significant figures to reflect analytical precision.

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-24-2014 Date of Receipt: 02-25-2014 Date of Report: 02-26-2014

MoldRANGETM: Extended Outdoor Comparison

Outdoor Location: 21402001-4TM20OUT

Fungi Identified	Outdoor		Typica	l Outo	loor Da	ıta for	:	Typical Outdoor Data for:					
	data	Feb	ruary ir	n Calif	ornia† (n‡=16	302)	The entire year in California† (n‡=200710)					
	spores/m3	very low	low	med	high	very high	freq %	very low	low	med	high	very high	freq %
Generally able to grow indoors*													
Alternaria	40	13	13	20	44	67	38	13	13	27	67	110	53
Bipolaris/Drechslera group	-	7	13	13	27	40	7	7	13	13	27	40	12
Chaetomium	-	7	13	13	27	40	9	8	13	13	27	47	19
Cladosporium	4,800	80	150	400	1,000	1,700	95	110	210	610	1,600	2,800	97
Curvularia	-	7	10	13	13	27	2	7	13	13	27	53	6
Nigrospora	-	7	13	13	13	27	4	7	13	13	27	53	8
Other brown	13	11	13	13	27	47	29	13	13	13	40	53	34
Penicillium/Aspergillus types	110	53	66	200	490	820	82	53	100	210	590	1,000	84
Stachybotrys	-	13	13	13	40	80	3	7	13	13	33	67	4
Stemphylium	13	7	13	13	27	40	5	7	13	13	27	40	9
Torula	80	7	13	13	40	53	5	8	13	13	40	67	12
Seldom found growing indoors**													
Ascospores	590	27	53	160	510	960	72	25	53	110	360	690	71
Basidiospores	1,300	53	110	430	1,800	3,900	95	53	80	260	990	2,300	93
Oidium	-	13	13	13	40	67	14	13	13	13	44	75	19
Rusts	53	8	13	13	40	73	14	13	13	13	53	80	26
Smuts, Periconia, Myxomycetes	80	13	13	27	67	110	54	13	13	40	110	210	68
§ TOTAL SPORES/m3	7,100												

[†]The 'Typical Outdoor Data' represents the typical outdoor spore levels for the location and time frame indicated. The last column represents the frequency of occurrence. The very low, low, med, high, and very high values represent the 10, 20, 50, 80, and 90 percentile values of the spore type when it is detected. For example, if the frequency of occurrence is 63% and the low value is 53, it would mean that the given spore type is detected 63% of the time and, when detected, 20% of the time it is present in levels above the detection limit and below 53 spores/m3. These values are updated periodically, and if enough data is not available to make a statistically meaningful assessment, it is indicated with a

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor data" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. In addition, EMLab P&K may not have received and tested a representative number of samples for every region or time period. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the use or interpretation of the data contained in, or any actions taken or omitted in reliance upon, this report.

EMLab P&K, LLC EMLab ID: 1175106, Page 1 of 1

[§] Total Spores/m3 has been rounded to two significant figures to reflect analytical precision.

^{*} The spores in this category are generally capable of growing on wet building materials in addition to growing outdoors. Building related growth is dependent upon the fungal type, moisture level, type of material, and other factors. *Cladosporium* is one of the predominant spore types worldwide and is frequently present in high numbers. *Penicillium/Aspergillus* species colonize both outdoor and indoor wet surfaces rapidly and are very easily dispersed. Other genera are usually present in lesser numbers.

^{**} These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens. However, in each group there are notable exceptions. For example, agents of wood decay are members of the basidiomycetes and high counts of a single morphological type of basidiospore on an inside sample should be considered significant.

 $[\]ddagger$ n = number of samples used to calculate data.

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-24-2014 Date of Receipt: 02-25-2014 Date of Report: 02-26-2014

MoldSTATTM: Supplementary Statistical Spore Trap Report

Outdoor Summary: 21402001-4TM20OUT:

Species detected		Outdoo	r sample s	pores/m3	Typical outdoor range	s Freq.
	<100	1K	10K	>100K	(North America)	%
Alternaria				40	7 - 33 - 590	45
Ascospores				590] 13 - 210 - 5,700	76
Basidiospores				1,300] 17 - 450 - 24,000	92
Cladosporium				4,800] 27 - 470 - 10,000	90
Other brown				13] 7 - 13 - 130	24
Penicillium/Aspergillus types				110] 13 - 170 - 2,700	68
Rusts				53] 7 - 20 - 360	20
Smuts, Periconia, Myxomycetes				80] 7 - 53 - 930	64
Stemphylium				13] 7 - 13 - 93	3
Torula				80] 7 - 13 - 190	9
Total				7,100		

The "Typical outdoor ranges" and "Freq. %" columns show the typical low, medium, and high spore counts per cubic meter and the frequency of occurrence for the given spore type. The low, medium, and high values represent the 2.5, 50, and 97.5 percentile values when the spore type is detected. For example, if the low value is 53 and the frequency of occurrence is 63%, it would mean that we typically detect the given spore type on 63 percent of all outdoor samples and, when detected, 2.5% of the time it is present in levels below 53 spores/m3.

Indoor Samples

Location: 21402001-4TM21

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: < 1%	dF: 5 Result: 6.2143 Critical value: 11.0705 Inside Similar: Yes	Result: 0.0000	dF: N/A Result: N/A Critical value: N/A Outside Similar: N/A	Score: 100 Result: Low
Species	Detected		Spores/m3	
		<100 1K	10K	>100K
	None Detected			

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: < 1%	dF: 5 Result: 6.2143 Critical value: 11.0705 Inside Similar: Yes	Result: 0.0000	dF: N/A Result: N/A Critical value: N/A Outside Similar: N/A	Score: 100 Result: Low
Species	Detected		Spores/m3	
		<100 1K	10K	>100K
	None Detected			< 13

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu Re: 21402001-4 Date of Sampling: 02-24-2014 Date of Receipt: 02-25-2014 Date of Report: 02-26-2014

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 21402001-4TM23

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)		
Result: 4%	dF: 5 Result: 6.2143 Critical value: 11.0705 Inside Similar: Yes	Result: 0.4615		dF: 10 Result: 0.5091 Critical value: 0.5515 Outside Similar: No	Score: 116 Result: Low		
Species 1	Detected			Spores/m3			
		<100	1K	10K	>100K		
	Basidiospores				210		
	Cladosporium				110		
	Other brown				13		
	Total				330		

Location: 21402001-4TM24

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)					
Result: 1%	dF: 5 Result: 6.2143 Critical value: 11.0705 Inside Similar: Yes	Result: 0.4286		dF: 11 Result: 0.0273 Critical value: 0.5273 Outside Similar: No	Score: 105 Result: Low					
Species 1	Detected	Spores/m3								
		<100	1K	10K	>100K					
	Cladosporium				53					
	Oidium				13					
	Other brown				13					
	Rusts				13					
	Total				93					

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)					
Result: 2%	dF: 5 Result: 6.2143 Critical value: 11.0705 Inside Similar: Yes	Result: 0.4615	dF: 10 Result: 0.6424 Critical value: 0.5515 Outside Similar: Yes	Score: 102 Result: Low					
Species	Detected	Spores/m3							
		<100 1K	10K	>100K					
	Basidiospores			53					
	Cladosporium			110					
	Rusts			13					
	Total			170					

Date of Sampling: 02-24-2014

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Date of Receipt: 02-25-2014 Re: 21402001-4 Date of Report: 02-26-2014

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 21402001-4TM26

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)				
Result: 1%	dF: 5 Result: 6.2143 Critical value: 11.0705 Inside Similar: Yes	Result: 0.3333	dF: 10 Result: 0.6273 Critical value: 0.5515 Outside Similar: Yes	Score: 108 Result: Low				
Species	Detected		Spores/m3					
		<100 1K	10K	>100K				
	Cladosporium			53				
Penic	illium/Aspergillus types			53				
	Total			110				

^{*} The Friedman chi-square statistic is a non-parametric test that examines variation in a set of data (in this case, all indoor spore counts). The null hypothesis (H0) being tested is that there is no meaningful difference in the data for all indoor locations. The alternative hypothesis (used if the test disproves the null hypothesis) is that there is a difference between the indoor locations. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

**** MoldSCORETM is a specialized method for examining air sampling data. It is a score between 100 and 300, with 100 indicating a greater likelihood that the airborne indoor spores originated from the outside, and 300 indicating a greater likelihood that they originated from an inside source. The Result displayed is based on the numeric score given and will be either Low, Medium, or High, indicating a low, medium, or high likelihood that the spores detected originated from an indoor source. EMLab P&Kreserves the right to, and may at anytime, modify or change the MoldScore algorithm without notice.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor ranges" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. With the statistical analysis provided, as with all statistical comparisons and analyses, false-positive and false-negative results can and do occur. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the data contained in, or any actions taken or omitted in reliance upon, this report.

EMLab ID: 1175106, Page 3 of 3 EMLab P&K, LLC

^{**} An agreement ratio is a simple method for assessing the similarity of two samples (in this case the indoor sample and the outdoor summary) based on the spore types present. A score of one indicates that the types detected in one location are the same as that in the other. A score of zero indicates that none of the types detected indoors are present outdoors. Typically, an agreement of 0.8 or higher is considered high.

^{***} The Spearman rank correlation is a non-parametric test that examines correlation between two sets of data (in this case the indoor location and the outdoor summary). The null hypothesis (H0) being tested is that the indoor and outdoor samples are unrelated. The alternative hypothesis (used if the test disproves the null hypothesis) is that the samples are similar. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Date of Sampling: 02-24-2014 Date of Receipt: 02-25-2014 Date of Report: 02-26-2014 Re: 21402001-4

 $\mathbf{MoldSCORE}^{\mathsf{TM}}\mathbf{:}\ \mathbf{Spore}\ \mathbf{Trap}\ \mathbf{Report}$ Outdoor Sample: 21402001-4TM20OUT

Fungi Identified	Ου	ıtd	00	r	san	ıpl	e	spo	re	s/ı	m3	Raw	Spores/
_	<100	0			1K			10K		>1	100K	count	m3
Generally able to grow indoors*													
Alternaria												3	40
Bipolaris/Drechslera group												ND	< 13
Chaetomium												ND	< 13
Cladosporium												90	4,800
Curvularia												ND	< 13
Nigrospora												ND	< 13
Other brown												1	13
Penicillium/Aspergillus types†												2	110
Stachybotrys												ND	< 13
Stemphylium												1	13
Torula												6	80
Seldom found growing indoors**													
Ascospores												11	590
Basidiospores												24	1,300
Rusts												4	53
Smuts, Periconia, Myxomycetes												6	80
Total													7,053

Fungi Identified	In	doo	r sa	mp	le s	spor	es/n	n3	Raw	Spores/
	<100	1	1 K			10K	>	100F	count	m3
Generally able to grow indoors*										
Alternaria									ND	< 13
Bipolaris/Drechslera group									ND	< 13
Chaetomium									ND	< 13
Cladosporium									ND	< 13
Curvularia									ND	< 13
Nigrospora									ND	< 13
Penicillium/Aspergillus types†									ND	< 13
Stachybotrys									ND	< 13
Torula									ND	< 13
Seldom found growing indoors**										
Ascospores									ND	< 13
Basidiospores									ND	< 13
Rusts									ND	< 13
Smuts, Periconia, Myxomycetes									ND	< 13
Total						·				N/A

100	MoldSCORE 100 200 300								
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
Fina	l MoldSC	ORE	100						

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-24-2014 Date of Receipt: 02-25-2014 Date of Report: 02-26-2014

MoldSCORETM: Spore Trap Report

Location: 21402001-4TM22

Fungi Identified	Indo	or	samj	ple s	spore	s/n	13	Raw	Spores/
	<100		1K		10K	>1	100K	count	m3
Generally able to grow indoors*									
Alternaria								ND	< 13
Bipolaris/Drechslera group								ND	< 13
Chaetomium								ND	< 13
Cladosporium								ND	< 13
Curvularia								ND	< 13
Nigrospora								ND	< 13
Penicillium/Aspergillus types†								ND	< 13
Stachybotrys								ND	< 13
Torula								ND	< 13
Seldom found growing indoors**									
Ascospores								ND	< 13
Basidiospores								ND	< 13
Rusts								ND	< 13
Smuts, Periconia, Myxomycetes								ND	< 13
Total			•						N/A

MoldSCORE:	
	100
	100
	100
	100
	100
	100
	100
	100
	100
	100
	100
	100
	100
Final MoldSCORE	100

Fungi Identified	In	Indoor sample spores/m3							5	Kaw	Spores/	
	<100	1		١K			10K		>10	0K	count	m3
Generally able to grow indoors*												
Alternaria											ND	< 13
Bipolaris/Drechslera group											ND	< 13
Chaetomium											ND	< 13
Cladosporium											2	110
Curvularia											ND	< 13
Nigrospora											ND	< 13
Other brown											1	13
Penicillium/Aspergillus types†											ND	< 13
Stachybotrys											ND	< 13
Torula											ND	< 13
Seldom found growing indoors**												
Ascospores											ND	< 13
Basidiospores											4	210
Rusts											ND	< 13
Smuts, Periconia, Myxomycetes											ND	< 13
Total												333

100	MoldSCORE; 200 300 Score									
			100							
			100							
			100							
			100							
			100							
			100							
			105							
			100							
			100							
			100							
			100							
			116							
			100							
			100							
Fina	Final MoldSCORE									

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-24-2014 Date of Receipt: 02-25-2014 Date of Report: 02-26-2014

MoldSCORETM: Spore Trap Report

Location: 21402001-4TM24

Fungi Identified	Inc	loor	sam	ple	sp	ore	es/n	13	Raw	Spores/
	<100		1K		1	0K	>	100K	count	m3
Generally able to grow indoors*										
Alternaria									ND	< 13
Bipolaris/Drechslera group									ND	< 13
Chaetomium									ND	< 13
Cladosporium									1	53
Curvularia									ND	< 13
Nigrospora									ND	< 13
Other brown									1	13
Penicillium/Aspergillus types†									ND	< 13
Stachybotrys									ND	< 13
Torula									ND	< 13
Seldom found growing indoors**										
Ascospores									ND	< 13
Basidiospores									ND	< 13
Oidium									1	13
Rusts									1	13
Smuts, Periconia, Myxomycetes				Ш			П		ND	< 13
Total										93

100	ORE;	Score	
			100
			100
			100
			100
			100
			100
			105
			100
			100
			100
			100
			100
			105
			105
			100
Fina	l MoldSC	ORE	105

Fungi Identified	In	Indoor sample spores/m3			Raw	Spores/				
	<100)		1K		10K	>10	0K	count	m3
Generally able to grow indoors*										
Alternaria							Ш		ND	< 13
Bipolaris/Drechslera group									ND	< 13
Chaetomium							Ш		ND	< 13
Cladosporium									2	110
Curvularia									ND	< 13
Nigrospora									ND	< 13
Penicillium/Aspergillus types†									ND	< 13
Stachybotrys									ND	< 13
Torula									ND	< 13
Seldom found growing indoors**										
Ascospores									ND	< 13
Basidiospores									1	53
Rusts									1	13
Smuts, Periconia, Myxomycetes									ND	< 13
Total										173

MoldSCORE‡					
100 200 300 Scor					
			100		
			100		
			100		
			100		
			100		
			100		
			100		
			100		
			100		
			100		
			102		
			105		
			100		
Fin	Final MoldSCORE 102				

Client: Hygiene Technologies International, Inc.

C/O: Mr. Lakhpreet Sandhu

Re: 21402001-4

Date of Sampling: 02-24-2014 Date of Receipt: 02-25-2014 Date of Report: 02-26-2014

MoldSCORETM: Spore Trap Report

Location: 21402001-4TM26

Fungi Identified	Indo	or sam	Raw	Spores/		
	<100	1K	10K	>100	K count	m3
Generally able to grow indoors*						
Alternaria					ND	< 13
Bipolaris/Drechslera group					ND	< 13
Chaetomium					ND	< 13
Cladosporium					1	53
Curvularia					ND	< 13
Nigrospora					ND	< 13
Penicillium/Aspergillus types†					1	53
Stachybotrys					ND	< 13
Torula					ND	< 13
Seldom found growing indoors**						
Ascospores					ND	< 13
Basidiospores					ND	< 13
Rusts					ND	< 13
Smuts, Periconia, Myxomycetes					ND	< 13
Total						107

MoldSCORE; 100 200 300 Score						
100 200	300	beore				
		100				
		100				
		100				
		100				
		100				
		100				
		108				
		100				
		100				
		100				
		100				
		100				
		100				
Final MoldSCC	RE	108				
·	·					

^{*} The spores in this category are generally capable of growing on wet building materials in addition to growing outdoors. Building related growth is dependent upon the fungal type, moisture level, type of material, and other factors. *Cladosporium* is one of the predominant spore types worldwide and is frequently present in high numbers. *Penicillium/Aspergillus* species colonize both outdoor and indoor wet surfaces rapidly and are very easily dispersed. Other genera are usually present in lesser numbers.

** These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens.

EMLab P&K, LLC EMLab ID: 1175106, Page 4 of 4

^{**} These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens. However, in each group there are notable exceptions. For example, agents of wood decay are members of the basidiomycetes and high counts of a single morphological type of basidiospore on an inside sample should be considered significant.

[†]The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods.

[‡]Rated on a scale from 100 to 300. A rating less than 150 is low and indicates a low probability of spores originating inside. A rating greater than 250 is high and indicates a high probability that the spores originated from inside, presumably from indoor mold growth. A rating between 150 and 250 indicates a moderate likelihood of indoor fungal growth. MoldSCORE is NOT intended for wall cavity samples. It is intended for ambient air samples in residences. Using the analysis on other samples (like wall cavity samples) will lead to misleading results.



Hygiene Technologies International, Inc.

₃₆₂₅ 001168914

(310) 370-8370 (310) 370-2474 FAX www.hyglenotech.com

	<u>+</u>				
Project Number/Purcha		21402	, <u> </u>		
Project Contact:	lomo e.	m/k.hs	Turnaround Required:		
Lab Destination:	EML	ABPBK	- Lab Contact: SAMPLE 12-6 COLVI		
SAMPLE ID	VOLUME	MEDIA	ANALYSIS REQUESTED		
21402001-4 TMO look	- 75L	Piracell			
21402001-4 TMO2	75L				
2140200)-4-13403					
21402007-4+MU4	75L				
21402001-4 TM05	751				
214021101-4-TM06	35L	1/			
		i			
			·		
Special Instructions:	Land	low so	mpling (Pa-1)		
		-			
·		912614			
1. Sampled by:	Endly or	14009:56	Received by: 166 10-00 02/01/14 1292pm		
2. Relinquished by: ≤	tom Dun o	~ 2/7/14@12	Received by:		
3. Relinquished by: Received by:					
		Please include signa	ture, date, and fine		
Lab Use Only:					
			•		
			i		
			I I		



Hygiene Technologies International, inc.

3625 Del Ar Tomanos,

(310) 370-8370

(310) 370-8370 (310) 370-2474 FAX www.hyglenetech.com

	<u> </u>				(ss)
Project Number/ Purcha			1-4-	Date Submitt	ed: 2/8/14
Project Contact: 6-8cm dlu / King Turnaround Required: No Smal					
Lab Destination:	<u>BMLAR</u>	PBIL_	Lab Contact:	Semp	le Receiving
SAMPLE ID	VOLUME	MEDIA		LYSIS REQU	
21402001-14-TMOTOUT	75L	A HED CELL	SPORE	TEGO	may si3 (18th)
21402001-4-TMOS	75L			l	
21402W14 TMO9	75.1				
21402001-4-TM10	75L			1	
21402VOI-47MII	_7 5 L				
21402001-4-TM12	<u> 75L</u>			V	
2140200-4-7m13	75L			1	
·					
		· · ·			
<u></u>					
<u>-</u>					
<u></u>					
The St. W.	<u>i</u>				
Special Instructions: _	Remisson	n Samo	Timy (R-2)	
· .			1	<u> </u>	
		(D .		
1. Sampled by:	tur Oly (m 2/14/401	Received by:	ll Wo	02/15/14 1230
2. Relinquished by:	Standbury U	n 2/19/14/01	70345		
3. Relinquished by:			Received by:		"
<u> </u>	<u>. I</u>	lease include signat	ure, date, and time		
Lab Use Only:					

Hygiene Technologies International, Inc.

3625 Del. 001173747 Torran

— — ক্ষেত্ৰস্থেত্যওস্থে (310) 370-2474 FAX www.hyglenetech.com

Project Number/Purcha	se Order: 2	(40200	-4 Date Submitted: 2/20/14
Project Contact:		1 .	Turnaround Required: Normal
Lab Destination:	EMI	A13 P8K	' . 1
SAMPLE ID	VOLUME	MEDIA	ANALYSIS REQUESTED
21402061-47M1400t	75L	An Kro-Cell	Spare Toap Anglysis (Total)
21402001-4 TMIS	75L		
21402001-4 TM16	75L	·	
21402001477417	75L		
2 402001-47M18	751		
21402001-4 TM19	7-5L	_ ~	
			· · · · · · · · · · · · · · · · · · ·
<u></u>	· · · · · · · · · · · · · · · · · · ·		
·			
·-····································			
allo VI Nato	<u></u> _		
Special Instructions: _	<u>Pen</u>	Don	Sampling (R-3)
	····		
	· · · · · · · · · · · · · · · · · · ·		
1. Sampled by:	andly or	~ 2/20/14 @	Received by: 2/21/14@08/
2. Relinquished by:	# mom	on 2/20/146	Received by:
3. Relinquished by: _			Received by:
	1	Please include signar	ure, date, and time
Lab Use Only:			
ζ			



Hygiene Technologies International, Inc.

3625 Dc 001175106 Tom

> — — roz 370-6370* (310) 370-2474 FAX www.hygienetech.com

Project Number/Purcha	se Order:	14020	01-4 Date Submitted: 2/24/14		
Project Contact:		1	· · · · · · · · · · · · · · · · · · ·		
Lab Destination:	12ML	AB PBK	Lab Contact: Sample Receiving		
SAMPLE ID	VOLUME	MEDIA	ANALYSIS REQUESTED		
21402001-4-Tm2001	675L	Amo-cell	Sporetrap AnalyBCH		
21402001-4 TM21	75L	Anto-Cell	, , , , , , , , , , , , , , , , , , , ,		
21402061-4 JM22	756	ATT-0-Cell			
21402001471423	751	Aix-o-Cell			
2140200147M24	75L	AIX-O-Cell			
214020014 TM25		Arro-Cell			
214/2001471126	75L_	Alxa-cell			
· · · · · · · · · · · · · · · · · · ·		<u> </u>			
			· · · · · · · · · · · · · · · · · · ·		
			·		
Special Instructions:	2 em	dum !	Sampling (P-4)		
			1 1 2		
1. Sampled by:	Emster in	1 2/24/146	Received by: Life Was Debs-hu 1255		
1. Sampled by: Family on 2/24/14 Received by: Let the 02/05/14 1255 2. Relinquished by: Family on 2/24/14 Received by:					
3. Relinquished by:	1	· · · · · · · · · · · · · · · · · · ·	m ,		
Flease include signature, date, and time					
Lab Use Only:					
			1		